

3.2

Geophysical risk: volcanic activity

Sue Loughlin, Sara Barsotti, Costanza Bonadonna, Eliza Calder

3.2.1 Volcanoes and volcanic activity

Volcanoes provide spectacular evidence of the dynamic nature of planet Earth and bring many long-term benefits to society, including rich soils, tourism and geothermal energy. Some erupt frequently and others may appear benign for generations, which means the risk they pose may be underestimated. Understanding the risk first requires characterisation of the volcano and knowledge of the type, magnitude and frequency of past eruptions.

3.2.1.1 Global distribution of volcanoes and volcanoes in Europe

There are about 1 550 known terrestrial volcanoes that have erupted in the past $\approx 10\,000$ years and are therefore likely to erupt again in the future;

they are described as ‘active’ (Siebert et al., 2010; Cottrell, 2014).

Volcanic eruptions may cause local to global impacts; in order to understand and mitigate risks, the first step is to recognise a volcano as active and to characterise its past activity.

Most have formed along colliding or diverging tectonic plate boundaries (e.g. the Pacific margins, the Mediterranean, the Lesser Antilles and Iceland; Figure 3.7) and these account for $>94\%$ of known historical eruptions (Siebert et al., 2015); the remainder have formed above mantle ‘hotspots’ (e.g. Hawaii).

In Europe, volcanism from Spain

(Bartolini et al., 2015) to Armenia (Savov et al., 2016) is mainly caused by the convergence of the northward-moving African and Arabian lithospheric plates with the Eurasian plate and microplates in the Aegean Sea and Anatolia (Figure 3.8). In Iceland, volcanism is caused by a combination of rifting at the Mid-Atlantic Ridge and a ‘hotspot’. There are 32 volcanoes in Iceland (Ilyinskaya et al., 2015), 47 known volcanoes in continental Europe (Siebert et al., 2010), and many more in autonomous regions, European dependencies and territories in the Atlantic (Canary Islands, Azores, Cabo Verde, Tristan da Cunha, Ascension Island), the Lesser Antilles (Montserrat, Guadeloupe, Martinique, Saba) and the Indian Ocean (La Réunion). About 15 million people in Europe live within just 30 km of an active volcano; of these, more than 2.2 million live within 20 km of the Campi Flegrei caldera in Italy and more than 675 000 live within 10 km of Vesuvius (Siebert et al., 2010).

3.2.1.2 Eruption type, duration, frequency and size

Globally, about 70 volcanoes erupt each year and at any one time at least 20 are erupting (Siebert et al., 2010, 2015). Eruptions are complex time-dependent events, which often exhibit distinct phases including effusive (e.g. lava flows/domes) and/or explosive types of activity (e.g. Gudmundsson et al., 2012) over durations

of hours to decades (Brown et al., 2015).

Major controls on eruption type include magma chemistry, rheology and volatile content. Eruptions can be measured using magnitude (erupted mass), but volume is often used as a proxy for magnitude for explosive eruptions (e.g. the Volcanic Explosivity Index, see Newhall and Self 1982, Pyle 2015).


Some volcanoes erupt frequently (e.g.

Stromboli and Etna), whereas others (e.g. Campi Flegrei) erupt infrequently, with hundreds of years between eruptions (e.g. Selva et al., 2012; Brown et al., 2014). Global data show a power law relationship between magnitude and frequency, such that larger magnitude eruptions are less frequent (Deligne et al., 2010). In order to understand the distribution of eruption types and magnitudes in time and space at a given volcano (and, therefore, the likelihood and type of future

FIGURE 3.7

The locations of known Holocene (past $\approx 10,000$ years) terrestrial volcanoes of the world, most of which form near tectonic plate boundaries (Bird, 2003).

Source: Smithsonian Institution (2013)



eruptions), geological and geochronological studies are an essential starting point (e.g. Druitt et al., 1999; Orsi et al., 2004; Thordarson and Larsen, 2007; Hicks et al., 2012).

3.2.1.3 Causes of volcanic unrest and eruptions

Eruptions are caused by complex processes including magma overpressurisation. Most eruptions are preceded by


one or multiple episodes of ‘volcanic unrest’ as magma moves towards the Earth’s surface (Acocella et al., 2015; Parks et al., 2015). The movement of magma through the crust (and its interaction with hydrothermal systems) causes pressure changes, which result in ground deformation and earthquakes, and also induces detectable changes in mass and/or density (Frey-mueller et al., 2015). During magma ascent, volatiles (gases) separate and are either retained in the magma as

bubbles or escape to interact with the hydrothermal system or be released at the surface. (Aiuppa et al., 2013). An episode of volcanic unrest may last for a matter of days to a number of years (average ≈ 500 days), and understanding the processes driving unrest and eruption is an essential part of effective early warning (Cashman et al. 2013, Sparks and Cashman 2017). Volcanoes that erupt infrequently (e.g. calderas) may experience many episodes of unrest (e.g. De Natale

FIGURE 3.8

Maximum known Volcanic Explosivity Index (0-8) of eruptions at European volcanoes in the past $\approx 10\,000$ years, based on the Smithsonian Institution Volcanoes of the World database (VOTW4.22). Volcanoes with unknown eruption histories are marked as black triangles.

Source: Smithsonian Institution (2013)



et al., 2006). In contrast, in a global study of 228 volcanoes active between 2000 and 2011 (many of which erupt frequently) the ‘Volcanic unrest in Europe and Latin America’ project (VUELCO) funded by the European Union’s 7th Framework Programme (FP7) showed that 47 % of documented periods of volcanic unrest led to an eruption (Phillipson et al., 2013). Some episodes of detected unrest may not be caused by magma and may be entirely tectonic or caused by hydrothermal phenomena (e.g. Segall, 2013; Biggs et al., 2009).

3.2.2 Monitoring systems and early warning

Volcano observatories are the official institutions in charge of monitoring volcanoes. They may be dedicated to a single volcano (e.g. the Montserrat Volcano Observatory) or may operate from national institutions and be responsible for multiple volcanoes in a country (e.g. the Icelandic Met Office and Istituto Nazionale di Geofisica e Vulcanologia). Some institutions have responsibility for volcanoes and volcano observatories overseas (e.g. Institut de Physique du Globe de Paris).

Volcano observatories have a key role in early warning. They collect multiple streams of diverse data, analyse the data in near real-time, determine the level of threat and make decisions on, for example, raising alert levels (Villagrán de León, 2012). These decisions must be based on sound evidence (Marzocchi et al., 2012; Bretton et al., 2015). The quality, range and sophistication of monitoring methods has increased dramatically in recent years

(Sparks et al., 2012), with advances in computing underpinning improvements in power, speed, data transmission, data analysis and modelling techniques. Long-term monitoring at quiescent volcanoes is necessary to establish baselines, and satellite remote sensing provides many opportunities as the spatial and temporal resolution of data improves (e.g. Harris et al., 2016; Bagnardi et al., 2016). Nevertheless, only a small fraction of the world’s 1 550 volcanoes have sufficient ground monitoring and the necessary accompanying institutional capacities to effectively support DRM, despite evidence that volcano monitoring is cost-effective (Newhall et al., 1997).

3.2.2.1 Geophysical monitoring (seismic, deformation, gas, infrasound) and the need for global monitoring

Episodes of unrest are highly variable in character, so forecasting the onset of an eruption remains a significant challenge (Chiodini et al., 2016; Selva et al., 2015; Marzocchi and Bebbington, 2012; Sigmundsson et al., 2010). Accelerating rates of seismicity and deformation may be detected before eruptions (Sigmundsson et al., 2010; Saltogianni et al., 2014; Cannavò et al., 2015) and tracking the location of volcano-tectonic earthquakes in near real time (Thorkelsson, 2012; Sigmundsson et al., 2015, Pallister and McNutt, 2015; Falsaperla and Neri, 2015) may facilitate eruption forecasting. Long-period earthquakes and micro-earthquakes can be key indicators of imminent eruption, especially during an ongoing eruption (McNutt et al., 2015). Cyclic patterns

of activity can also enable forecasting of hazardous events (Voight et al., 1999; Loughlin et al., 2002). Borehole strainmeters have successfully been used to forecast eruptions of Hekla in Iceland (Roberts et al., 2011).

If appropriate monitoring is in place at a volcano, it may be possible to issue short-term forecasts of eruptions and volcanic activity and to provide early warnings for different hazards.

Although satellite passes are not yet frequent enough to use Interferometric Synthetic Aperture Radar (InSAR) as a forecasting tool, it can be used in combination with other data to gain tremendous insights into volcanic unrest and eruption (e.g. Gudmundsson et al., 2016; Spaans and Hooper, 2016). InSAR is useful to detect deformation at remote volcanoes and at regional scales (Biggs et al., 2014; Parks et al., 2015).

Gas emissions (Silva et al., 2015; Aiuppa et al., 2013; Chiodini et al., 2015), the chemistry, temperature and level of crater lakes and groundwater (Hernández et al., 2007), and the geochemistry and flow rates of glacial rivers (Kristmansdóttir et al., 1999) may all show detectable changes before and during eruptions. Gas emissions can be monitored using ground-based, airborne or satellite remote sensing (Aiuppa et al., 2007, 2010; Nadeau et al., 2011; Conde et al., 2013). The gas

most easily detected and monitored in the atmosphere during eruptions is sulphur dioxide (SO₂) (Oppenheimer et al., 2013; Flower et al., 2016).

Petrology and geochemistry can be used in near real time to characterise eruptive materials and understand magmatic properties and dynamics (Hartley et al., 2016; Pankhurst et al., 2014). Rapid analysis of tephra can detect whether or not there is a magmatic component to phreatic (steam-driven) eruptions (Suzuki et al., 2013).

Environmental monitoring such as dissolved constituents in rainwater, ash leachates (Witham et al., 2005) and particulate (air quality) monitoring can potentially provide information about both eruptive behaviour and probable impacts on health, the environment, infrastructure and buildings (Gislason et al., 2015).

3.2.2.2 Additional and emerging monitoring methods

Volcanic infrasound is a technique that detects, locates and characterises shallow or aerial acoustic sources at volcanoes (Fee and Matoza, 2013; Ulivieri et al., 2013). During the H2020 Atmospheric dynamics Research InfraStructure in Europe 2 (ARISE2) project, episodes of lava fountaining at Etna were recorded ≈ 600 km away, providing evidence that near-real-time notification of ongoing volcanic activity at a regional scale can be achieved (Johnson and Ripepe, 2011; Marchetti et al., 2016).

Establishing mass eruption rate (a parameter needed to effectively forecast

ash dispersal) and characterising ash clouds in near real time is a current challenge (Ripepe et al., 2013; Lamb et al., 2015; Marzano et al., 2013, 2016). Monitoring the extrusion rate of lava is crucial to anticipate the evolution of active lava flow fields or stability of lava domes. Time series digital elevation models (DEMs) collected by satellite at Merapi volcano in 2010 (through the International Space Charter), combined with ground monitoring, enabled increasing extrusion rates to be identified, leading to a rise in alert level and timely evacuations that saved thousands of lives (Surono et al., 2012; Pallister and Surono, 2015). Extrusion rates can be established from the ground, unmanned aerial vehicles or aircraft using a variety of methods (e.g. Wadge et al. 2014a, 2014b; Harris et al., 2005).

Characterisation of heat sources (Figure 3.9) during volcanic unrest and eruption can support scientific understanding of eruptive behaviour and timely response (Harris et al., 2016). During the 2014-15 Bárðarbunga eruption in Iceland, the Middle Infra-Red Observation of Volcanic Activity (MIROVA) system (Coppola et al., 2015) was used to chart the evolution of the eruption when access was limited and visibility was poor.

In seismology, deterministic eruption forecasting based on the failure forecast method (FFM) is showing potential (Boué et al., 2016).

Time series observations of volcanoes and their emissions using static or video cameras can yield important insights into hazardous processes. Citizen science, including community

FIGURE 3.9

Measuring the temperature of pyroclastic flow deposits in Montserrat (block and ash flow deposits).
Source and Copyright BGS/Government of Montserrat



monitoring, can fill observational and information gaps, raise awareness of hazards and risk, and engage communities at-risk (e.g. Stevenson et al., 2013; Stone et al., 2014; Wallace et al., 2015). During the 2014-15 eruption at Bárðarbunga volcano, Iceland, people could document their experiences of poor air quality due to the gas-rich eruption online (IMO, n.d).

WOVodat is a searchable, web-accessible global relational database containing time-series monitoring data from more than 100 eruption episodes; this will allow global trends in unrest and eruption data to be interrogated to assist forecasting at individual volcanoes (Venezky and Newhall,

2007; Widiwijayanti et al., 2015).

3.2.2.3 Communication, reporting and alert levels

During unrest or eruption, scientists communicate in a variety of ways (reports, forecasts, alert levels) using a variety of media (email lists, short message service (SMS), social media, television and radio) to suit the needs of information users (Solana et al., 2008; Haynes et al., 2008a; Mothes et al., 2015). Such users are diverse and include civil aviation authorities, civil protection authorities, businesses, tourist operators, the media and the public. Ideally, the content and

format of such communications are tailored to users' needs (e.g. Lechner et al., 2017; Doyle et al., 2014) and users have considered in advance their thresholds for action (e.g. Marzocchi et al., 2012; Hicks et al., 2014). During an emergency, joint formal reports can be particularly effective if scientists and civil protection authorities work well together, and if the content and format has been designed specifically with users in mind (e.g. Scientific advisory board of the Icelandic Civil Protection, 2015).

A volcano Early Warning System (EWS) requires that monitoring data are collected and interpreted by scientists, the level of threat is determined

FIGURE 3.10

Summary of alert levels and civil protection system response for Vesuvius volcano, Italy. Alert levels are established by INGV Vesuvio based on changing monitoring parameters. The civil protection system responds in each operative phase according to the alert level and the emergency plan.

Source: authors

ALERT LEVEL	STATE OF THE VOLCANO	ERUPTION PROBABILITY	TIME OF ERUPTION	OPERATIVE PHASE
Base	No significant variation of monitored parameters	Very low	Undefined	
Caution	Significant change of monitored parameters	Low	Indefinite, or not less than several months	I Caution
Warning	Further significant change in monitored parameters	Medium	From months to weeks	II Warning
Alarm	Appearance of phenomena and/or evolution of monitored parameters suggesting a pre-eruption dynamic	High	From weeks to days	III Alarm

Operative Phase I: Verification of contingency plans, constant contact between scientists and civil protection, checking of functionality and immediate availability of resources, infrastructure and services needed for subsequent alert levels.

Operative Phase II: Voluntary evacuation of red zone to alternative accommodation outside the zone of risk. All involved in the emergency plan alert and prepared for Phase III.

Operative Phase III: Evacuation of the red zone within three days.

and a decision to alert stakeholders is made (Fearnley, 2013). Some volcano observatories use Volcanic Alert Levels (VALs) to communicate changes in the status of volcanic activity that imply a changing probability of eruption (Gardner and Guffanti, 2006; Fearnley, 2013; Winson et al., 2014) or changing types of hazard (Potter et al., 2014). Notification of a change in VAL is usually accompanied by situation-specific information in the form of a more detailed report. VALs are developed to suit local situations and, as such, they vary worldwide. Some focus on unrest and eruption forecasting (Figure 3.10) and others acknowledge the changing phenomena and hazards of long-lived eruptions (Potter et al., 2014). In situations in which major and costly mitigation actions are triggered by volcanic EWSs (e.g. the evacuation of urban areas), quantitative, objective and rational scientific decision-making is essential to avoid accusations of ‘false alarms’ (see Chapter 3.2.3, Hincks et al., 2014).

The global network of nine Volcanic Ash Advisory Centres (VAACs) was set up by the International Civil Aviation Organization (ICAO) following aircraft encounters with ash clouds in the 1980s (Guffanti et al., 2010). Volcano observatories provide reports to VAACs to support the initiation of ash dispersal models and have the option to set an ‘aviation colour code’ representing the status of volcanoes in the context of likelihood of eruption and potential for ash emissions (Lechner et al., 2017). This system can run in parallel to a VAL system.

In situations on the ground in which it is acknowledged that there may be

little time for response, alerts may be sent out to authorities and the public via SMS, telephone, radio or social media (e.g. IMO, 2016; Stone et al., 2014; Mothes et al., 2015). EWSs for lahars and jökulhlaups (glacier floods in Iceland) are variable in terms of components but, in general, require monitoring (e.g. acoustic flow monitors) to detect flows in proximal environments that can alert authorities to sound sirens downstream so that communities can be evacuated.

3.2.3 Volcanic hazard assessment

Volcanic hazards are diverse and they can occur in different combinations and interact in different ways throughout the unrest, eruption and post-eruption period.

Volcanoes generate multiple hazardous processes, the short- and long-term forecasting of which involve diverse methods to anticipate hazard footprints in order to enable anticipation and mitigation of impacts.

Scientists are improving their ability to assess and forecast these hazards, their likely ‘footprints’, interactions and impacts over different timescales. Short-term and long-term forecasts, to support crisis response and planning, respectively, are based on a vari-

ety of different approaches depending primarily on data availability. Deterministic and probabilistic approaches to hazard are used and are appropriate in different circumstances. .

3.2.3.1 Hazard forecasting

Short-term forecasts can enable communities across broad areas to prepare for imminent hazards and impacts. For example, simple simulations of expected atmospheric dispersal and deposition of volcanic tephra based on monitoring/observation parameters, can be made available (e.g. for Etna at INGV (n.d.) and for Mount St Helens at USGS (2015), Hasegawa et al., 2015). Similarly, short-term dispersal forecasts of SO₂ (which may adversely affect human and live-stock health) can enable mitigation actions to be taken (e.g. Gislason et al., 2015). Such forecasts can also be achieved for lava flows and lahars, and, in some places, mitigation of lava flow impacts has been attempted using engineering measures.

Volcanic hazard process models still need further development to better simulate key processes; this is especially true for pyroclastic flows, surges and lahars, the assessment of which currently lags behind that of tephra dispersal and fall. The ability to model interacting hazards is also important, such as rainfall-triggered lahars (Jones et al., 2015), eruption column collapse into pyroclastic flows or pyroclastic flows into lahars.

Long-term volcanic hazard assessment is primarily based on characterising the past eruptive activity of a volcanic system and understanding

the recurrence rates of eruptions and the range of possibilities for future eruptions. Such assessments are often presented as hazard maps. Ideally, geological and historical studies are needed to establish eruption histories but sometimes such information is not available, further fieldwork is needed, or data simply doesn't exist. For example, fine-grained deposits (e.g. ash fall, surges, lateral blasts)

may be missing from the geological record, so thorough consideration of knowledge gaps and uncertainties is paramount in any hazard analysis (e.g. Engwell et al., 2013; Sparks et al., 2013, Bonadonna et al. 2012, 2015). Volcanologists can study analogue volcanoes and global databases to address knowledge gaps (e.g. Ogburn et al., 2015), or use methods such as expert elicitation in order to consider

and quantify uncertainties (Aspinall, 2006, 2010). Uncertainty should be acknowledged in all scientific decision-making, forecasts and assessments.

3.2.3.2 Volcanic hazard maps

Volcanic hazard maps can communicate information about one or a range of hazards including lahars, pyroclastic flows and surges, tephra fall (Macedonio et al., 2008), ballistics, lava flows (Richter et al., 2016), and, sometimes, less frequent hazards such as debris avalanches and monogenetic eruptions. An International Association of Volcanology and Chemistry of the Earth's Interior (IAVCEI) working group is reviewing current global practice (>200 published hazard maps). They defined five major classes of hazard map and found that >60 % of maps are based primarily on the geological history of the volcano (Figure 3.11), despite incomplete eruption histories that do not represent all past and possible future scenarios. Furthermore, >83 % of hazard maps use a qualitative 'high-medium-low' description to indicate likelihood of impact, but the meanings behind these terms are open to broad interpretation (Calder et al., 2015).


The IAVCEI working group have established that there is no single approach that suits all situations; different approaches may be suitable for different needs. Nevertheless, there is consensus that quantitative, accountable and defensible hazard maps are increasingly needed.

The group aims to collectively define

FIGURE 3.11

Synthetic examples (not a real volcano) of the appearance of five hazard map types found during the review:
 (a) geology-based map,
 (b) integrated qualitative map,
 (c) administrative map,
 (d) modelling-based map and
 (e) probabilistic map.

Copyright: Cambridge University Press



good practices. Scientific priorities to enhance hazard maps include (1) improved methods for probabilistic analysis, especially for lahar, pyroclastic flows and surges, (2) establishing methods to undertake hazard assessments for data-poor volcanoes and (3) approaches for multihazard, multisenario probabilistic modelling (Calder et al., 2015). Although probabilistic volcanic hazard maps exist for a few of the world's best studied volcanoes, they are far from being the norm.

Haynes et al. (2007) recognised that maps are rarely well understood by users and that three-dimensional (3D) visualisation can significantly help understanding.

3.2.3.3 Probabilistic volcanic hazard assessment

A variety of methods are used, often in combination, to generate probabilistic volcanic hazard assessments over different timescales.

For example, statistical methods can be used to assess recurrence rates and locations of vents, which, when combined with numerical simulations of volcanic processes (e.g. lava flow emplacement, tephra dispersal and fall), can create hazard curves for specific locations or hazard maps for larger areas (Connor et al., 2015). There may be high uncertainty in vent location, particularly if a volcano has vents distributed across its flanks or the area is a volcanic field comprising multiple eruption centres (Connor et al., 2012; Bebbington and Cronin, 2010).

Because of the variety and potential complexity of volcanic hazards,

probabilistic hazard maps commonly attempt to communicate information about only a single hazard at a time (Figure 3.11e). For example, volcanic flows are typically displayed as the spatial variation of inundation probability over a given period of time. Tephra fall might be assessed using contours of probability given a hazardous threshold of tephra thickness (Jenkins et al., 2015a) or contours of tephra thickness given a certain probability in order to better assess the associated impact (e.g. Bonadonna, 2006; Biass et al. 2014, 2016a). Probabilistic hazards assessments can be local to global in scale.

The first probabilistic assessment of global tephra fall hazard has been attempted for the 2015 United Nations International Strategy for Disaster Reduction (UNISDR) Global Assessment Report, based on a method developed for the regional scale (Jenkins et al., 2012, 2015a).

Once volcanic unrest begins there are multiple potential eruptive outcomes (scenarios) due to the dynamic complexity of volcanic systems. In this situation, probabilistic methods provide a basis for scientists to explore those outcomes, allocate probability estimates to them (Marzocchi et al., 2012; Selva et al., 2010) and communicate them to authorities to support rational decision-making (e.g. Sparks 2003, Marzocchi et al., 2007). The results can be tested using statistical procedures and allow comparisons between volcanoes and other natural and non-natural hazards (e.g. Scandone et al., 1993; Bayarri et al., 2015).

Current probabilistic approaches build on the idea of event trees

(Newhall and Hoblitt, 2002) and on Bayesian statistics (e.g. Papadopoulos and Orfanogianaki, 2005). The probability estimates allocated to each outcome/scenario might be empirical, or be based on expert discussion and elicitation, numerical simulations or a combination of methods (e.g. Aspinall, 2006, 2010; Marzocchi and Bebbington, 2012). These methods are also useful if applied regularly at long-lived or frequently active volcanoes where probabilities change and assessments can be compared over time (Pallister et al., 2010; Wadge and Aspinall, 2014). Similar approaches have now been applied at Vesuvius (Neri, 2008), Teide-Pico Viejo, Tenerife (Martí et al., 2008), and Auckland Volcanic Fields, New Zealand (Lindsay et al., 2010). The same principle has also been developed to generate tools (e.g. Marzocchi et al., 2008).

3.2.4 Volcanic risk assessment and mitigation

3.2.4.1 Vulnerability and exposure

Vulnerability is complex, dynamic and spatially variable with many facets including systemic, social, functional and economic vulnerability (e.g. Enhancing resilience of communities and territories facing natural and na-tech hazards (ENSURE) project, Menoni et al., 2012). Exposure contributes to vulnerability (Cutter 2013) and includes the people and assets exposed to the hazards. Volcanic unrest and eruptions tend to unfold over weeks to years, thereby enhancing the dynamic complexity of factors that

contribute to vulnerability and exposure (Galderisi et al., 2013; Zuccaro et al., 2014).

Volcanic tephra fall is the hazard that most frequently affects large populations and assets and has reasonable vulnerability estimates in risk models (Spence et al., 2005; Jenkins et al., 2015). Jenkins et al. (2014), as part of the Mitigate and Assess risk from Volcanic Impact on Terrain and human Activities (MIA-VITA) project, developed guidelines and methodologies for carrying out initial physical vulnerability assessments. This built on previous projects including the SPeeD project at Vesuvius and the EXPLORIS project (e.g. Zuccaro et al., 2008; Marti et al., 2008). Jenkins et al. (2015) categorised tephra fall impacts by sector and considered the relationship between hazard intensity (in that case ash thickness) and damage or disruption to each sector (buildings, critical infrastructure, agriculture). More data need to be collected to inform estimates of physical vulnerability of buildings and infrastructure, through: (1) collection of post-eruption damage data (e.g. Baxter et al., 2005; Wilson et al., 2011; Biass et al., 2016a, 2016b; Charbonnier et al., 2013; Jenkins et al., 2015b), (2) experimental testing of materials failure, or (3) using theoretical calculations of material strengths (e.g. Jenkins et al., 2014). Damage data and experimental data remain sparse so theoretical calculations can contribute to the development of vulnerability functions, which provide the probability of a certain level of damage as a function of hazard intensity (Jenkins et al., 2014; see also Chapter 2.4).

A dynamic pressure scale for building

FIGURE 3.12

Soufrière Hills Volcano and Plymouth, the capital of Montserrat, in October 1997.

Copyright: British Geological Survey/Government of Montserrat



damage by pyroclastic surges (Baxter et al., 2005), developed after experiences in Montserrat (Figure 3.12) and at Mount St Helens, has contributed to simulation work at Vesuvius and other European volcanoes in the EXPLORIS project (Baxter et al., 2008).

Studies on social vulnerability in volcanic risk are increasing in number and showing the value of semi-quantitative and qualitative assessments (e.g. Sword-Daniels, 2011; Sword-Daniels et al., 2014). Hicks and Few (2015) showed that during long-term eruptions, coping capacity, maintenance of well-being, recovery of losses and rebuilding of livelihoods are highly variable within populations and tend to be linked to preceding socio-economic conditions (Birkman, 2007). Socio-economic impacts are most likely to be experienced by those with pre-existing and inter-related sociocultural, political and economic vulnerabilities (Wisner et al., 2012; Gaillard 2008). Volcanic activity can have a disproportionate effect on livelihoods and economy because of high systemic vulnerability (Wilson et al., 2011, 2014; Jenkins et al., 2015a).

Comprehensive quantitative assessment of the impacts of all types of volcanic hazard is relatively new but is most advanced for tephra fall (Craig et al., 2016; Elissondo et al., 2016; Wilson et al., 2011, 2012, 2014; Magill et al., 2013). Socioeconomic impacts due to tephra fall are most likely to be documented in long-lived eruptions (e.g. Sword-Daniels, 2011; Sword-Daniels et al., 2014).

3.2.4.2 Risk assessment methodologies

Volcanic risk assessment is not as advanced as assessment of other hazards such as flooding, earthquakes and tropical cyclones. For the long-lasting eruption at Soufriere Hills Volcano, Montserrat, volcanic risk has been assessed in a regular and consistent way for 20 years (Aspinall et al., 2002; Aspinall, 2006; Wadge and Aspinall, 2014). After deriving event scenario probabilities and their uncertainties by elicitation, risks and uncertainties are quantified using Monte Carlo modelling, and the risk is presented as (1) societal risk expressed quantitatively as a curve of the probability of exceeding a given number of fatalities, (2) individual risk given as an annualised probability of death (from the volcano) for any person living in a specific area and (3) occupational risk given for people working under certain conditions in specific areas.

An example of a volcanic risk model is the KazanRisk loss model (risk-frontiers.com/kazanrisk.htm), which uses numerical dispersal modelling of ash fall in Greater Tokyo to estimate potential losses associated with building damage, clean-up and reductions in agricultural productivity. At regional to global scales, the CAPRA risk modelling platform (ecapra.org) has been used to provide preliminary estimates of potential building damage around active volcanoes in the Asia-Pacific Region using simplified volcanic hazard outputs from a statistical emulator (Jenkins et al., 2015a). Because some crucial aspects of vulnerability must be assessed qualita-

tively, there is a need to find innovative ways to integrate qualitative with quantitative data to assess volcanic risk (Hicks and Few, 2015). Novel interdisciplinary approaches are now being developed (e.g. STREVA project) that combine volcanological techniques, probabilistic decision support and social science methods to ensure that the benefits of even uncertain and incomplete knowledge are acted upon to reduce risk (e.g. Hicks et al., 2014; Barclay et al., 2014). Stirling (2010) highlighted that different analytical methods suit different epistemic conditions and acknowledging the state of knowledge is a good start in enabling effective risk analysis and communication..

'Forensic analysis' of past disasters provides a strong basis for learning (e.g. Voight 1990, Loughlin et al. 2002, Thordarson and Self, 2003; Bird et al., 2010; Ragona et al., 2011), and longitudinal studies can reveal valuable insights into causal processes behind impacts and disasters (e.g. Integrated Research on Disaster Risk FORIN project 2011). Such approaches are also being applied to understand recovery processes that are complex and can last for decades (Sword-Daniels et al., 2015).

3.2.4.3 Civil Protection, scientists and risk management

Volcano observatories and civil protection authorities, working together as well as with the public, have reduced fatalities due to volcanic activity worldwide. At least 50 000 lives were saved in the 20th century (Auker et al., 2013) and even more have been saved since 1985 (Voight et al., 2013).

Mutual understanding and trust develop with an investment of time and effort (Haynes et al., 2008a, 2008b) and it is too late to start when an emergency begins. Effective communication and decision-making during a rapidly changing emergency situation (Fischhoff, 2013; Doyle et al., 2014) will be facilitated by good planning, preparation and response protocols (Doyle et al., 2015; Bretton et al., 2015). Interdisciplinary and transdisciplinary approaches can bring a wide range of methods and experiences together (e.g. communities, scientists, authorities) to facilitate better understanding, analysis and communication of hazard and risk (Hicks et al., 2014; Barclay et al., 2014, 2015).

Volcanic unrest and eruptions can be

prolonged, which may cause disruption and have long-term socio-economic impacts. Tephra fall can cause damage and disruption across sectors and has potential health impacts (Horwell and Baxter, 2006; Carlsen et al., 2012); therefore, planning for clean-up and recovery is essential (Hayes et al., 2017).

Preparedness for volcanic unrest and eruption often takes the form of contingency plans, which can be practised (Figure 3.13) by scientists, authorities and other stakeholders, including the public (Hicks et al., 2014). Different types of exercises have been reported around the world (Figure 3.13), ranging from the training of small groups to international reaction-chain exercises (Lindsay et al., 2010; Ricci et al.,

2013). In high-risk urban settings (e.g. Naples), there are significant costs to mitigation actions, even to exercises, so direct and indirect costs and benefits need to be carefully considered to support decision-making (Marzocchi et al., 2012; Woo, 2014).

The 2010 eruption of Eyjafjallajökull volcano in Iceland demonstrated that even small eruptions can have global impacts (Ragona et al., 2011). Therefore, international collaboration is essential to ensure that lessons learned and scientific progress are translated into planning and preparation across all sectors (Schmidt et al., 2011, 2015; Bonadonna et al., 2012).

3.2.5 Conclusions and key messages

Partnership

Long-term collaboration and effective partnerships between scientists (operational and research) and civil protection authorities are particularly important for effective evidence-based risk management and emergency response. The recent FUTUREVOLC and MEDSUV projects (FP7) showed how Europe-wide research partnerships can support such national and Europe-wide DRM efforts in particular. Engagement with users of scientific and civil protection advice can improve the format and content of outputs, enhancing understanding, uptake and effective decision-making at all levels. The knowledge and experience of those at risk is increasingly recognised as important and their involvement in the design and development of DRM strategies can be highly effective.

FIGURE 3.13

An evacuation exercise for the entire population of Tristan da Cunha, South Atlantic.

Source: photograph courtesy of Anna Hicks



Knowledge

Hazard, impact, vulnerability, loss and recovery data are sparse in volcanology but are needed to produce better hazard and risk assessments. Detailed study of all future eruptions and their impacts is needed. Despite an overall need for increased quantification in volcanic risk, interdisciplinary collaboration is recommended to capitalise on both quantitative and qualitative approaches to risk, particularly in situations in which data are scarce. Progress in process understanding is needed to enable better anticipation of hazardous events. Frameworks for the optimal combination of hazard and vulnerability analysis across multiple temporal and spatial scales is needed for comprehensive risk assessments and proactive policies of risk reduction.

Innovation

There is an ongoing need for development in monitoring techniques, integrated analysis of ground and space data, hazard and vulnerability assessment methodologies and interdisciplinary/transdisciplinary science. A next important step for the volcanology community as a whole is to enhance innovation in hazard and risk assessment strategies. There is an increasingly urgent need for near-real-time global monitoring and a reporting platform to support the anticipation of volcanic events that have wide-reaching or humanitarian impacts. This will require collaborative approaches and innovative integration of data in a wide variety of formats and at different spatial and temporal resolutions.

REFERENCES CHAPTER 3 - SECTION I

Introduction

UNISDR, 2015. Sendai framework for disaster risk reduction 2015–2030. United Nations International Strategy for Disaster Reduction. http://www.wcdr.org/uploads/Sendai_Framework_for_Disaster_Risk_Reduction_2015-2030.pdf, [accessed 04 April 2016].

Woo G., 2010. Operational earthquake forecasting and risk management. *Seismological Research Letters* 81(5).

3.1 Geophysical risk: earthquakes

Akkar, S., Sucuoglu, H., Yakut, A., 2005. Displacement-based fragility functions for low- and mid-rise ordinary concrete buildings. *Earthquake Spectra* 21(4), 901-927.

Albini, P., Musson, R., Rovida, A., Locati, M., Gomez Capera, A., Vigano, D., 2014. The global earthquake history. *Earthquake Spectra* 30(2), 607–24.

Alcik, H., Ozel, O., Apaydin, N., Erdik, M., 2009. A study on warning algorithms for Istanbul earthquake early warning system. *Geophysical Research Letters* 36(5), L00B05.

Anhorn, J., Khazai, B., 2015. Open space suitability analysis for emergency shelter after an earthquake. *Natural Hazards and Earth System Sciences* 1 (2), 4263-4297.

Ansal, A., Akinci, A., Cultrera, G., Erdik, M., Pessina, V., Tonuk, G., Ameri, G., 2009. Loss estimation in Istanbul based on deterministic earthquake scenarios of the Marmara Sea region (Turkey). *Soil Dynamics and Earthquake Engineering* 29, 699–709.

ARISTOTLE, n.d. All Risk Integrated System Towards Trans-boundary holistic Early-warning. <http://aristotle.ingv.it>, [accessed 13 April, 2017].

ATC-13, 1985. Earthquake damage evaluation data for California. Applied Technology Council. Redwood City, CA, USA.

Basili, R., Tiberti, M.M., Kastelic, V., Romano, F., Piatanesi, A., Selva, J., Lorito, S., 2013. Integrating geologic fault data into tsunami hazard studies. *Natural Hazards Earth System Sciences* 13, 1025–1050.

Bazzurro, P., Cornell, C., 1999. Disaggregation of seismic hazard. *Bulletin of the Seismological Society of America* 89: 501-520.

Bendimerad, F., 2001. Loss estimation: a powerful tool for risk assessment and mitigation. *Soil Dynamics and Earthquake Engineering* 21 (5), 467-472.

Bilham, R., 2004. Earthquakes in India and the Himalaya: tectonics, geodesy and history. *Annals of Geophysics* 47 (2–3), 839–858.

Bird, P., 2003. An updated digital model of plate boundaries. *Geochemistry, Geophysics, Geosystems* 4 (3), 1027.

Bommer, J., Abrahamson, N., 2006. Why do modern probabilistic seismic-hazard analyses often lead to increased hazard estimates?. *Bulletin of the Seismological Society of America* 96 (6), 1967-1977.

Bommer, J., Douglas, J., Scherbaum, F., Cotton, F., Bungum, H., Fah, D., 2010. On the selection of ground-motion prediction equations for seismic hazard analysis. *Seismological Research Letters* 81, 783–793.

Bommer, J., Spence, R., Erdik, M., Tabuchi, S., Aydinoglu, N., Booth, E., Re, D., Pterken, D., 2002. Development of an Earthquake Loss Model for Turkish Catastrophe Insurance. *Journal of Seismology* 6, 431-446.

Brzev, S., Scawthorn, C., Charleson, A., Allen, L., Greene, M., Jaiswal, K., Silva, V., 2013. GEM Building Taxonomy Version 2.0, GEM Technical Report 2013-02 V1.0.0. GEM Foundation, Pavia, Italy. doi: 10.13117/GEM.EXP-MOD.TR2013.02.

Burton, C., Silva, V., 2015. Assessing Integrated Earthquake Risk in OpenQuake with an Application to Mainland Portugal Earthquake Spectra 32 (3), 1383-1403.

Calvi, G., Pinho, R., 2004. LESSLOSS — A European integrated project on risk mitigation for earthquakes and landslides. IUSS Press, Pavia, Italy.

Carreño, L., Cardona, O., Barbat, A., 2007. Urban seismic risk evaluation: a holistic approach *Natural Hazards*, 40, 137-172.

CIESIN, 2004. Global Rural-Urban Mapping Project (GRUMP) — Urban extents. Center for International Earth Science Information Network. <http://sedac.ciesin.columbia.edu/data/set/grump-v1-urban-extents> [Accessed 21 September, 2016].

Cornell, C., 1968. Engineering seismic risk analysis. *Bulletin of the Seismological Society of America* 58, 1583-1606.

Cotton, F., Scherbaum, F., Bommer, J., Bungum, H., 2006. Criteria for selecting and adjusting ground-motion models for specific target regions: application to central Europe and rock sites. *Journal of Seismology* 10, 137–156.

Crowley, H., Colombi, M., Silva, V., 2014. Epistemic uncertainty in fragility functions for European RC buildings. In: Ptilakis, K., Crowley, H., Kaynia, A.M. (Eds), SYNER-G: Typology definition and fragility functions for physical elements at seismic risk: buildings, lifelines, transportation networks and critical facilities, Springer, Dordrech, Netherlands, pp. 95-109.

Crowley, H., Miriam, C., Borzi, B., Faravelli, M., Onida, M., Lopez, M., Polli, D., Meroni, F., 2009. A comparison of seismic risk maps for Italy. *Bulletin of Earthquake Engineering* 7 (1), 149-180.

Crowley, H., Ozecebe, S., Spence, R., Foulser-Piggott, R., Erdik, M., Alten, K., 2012. Development of a European building inventory database. In: Proceedings of the 15th World Conference on Earthquake Engineering, Lisbon, Portugal.

D'Ayala, D., Meslem, A., Vamvatsikos, D., Porter, K., Rossetto, T., 2015. Guidelines for analytical vulnerability assessment of low/ mid-rise buildings, GEM Technical Report 2014-12. GEM Foundation, Pavia, Italy.

Danciu, L., Kale, O., Akkar, S., 2016. The 2014 Earthquake Model of the Middle East: ground motion model and uncertainties. *Bulletin of Earthquake Engineering*.

Delavaud, E., Cotton, F., Akkar, S., Scherbaum, F., Danciu, L., Beauval, C., Drouet, S., Douglas, J., Basili, R., Sandikkaya, M.A., Segou, M., Faccioli, E., Theodoulidis, N., 2012. Toward a ground-motion logic tree for probabilistic seismic hazard assessment in Europe. *Journal of Seismology* 16, 451–473.

Dolce, M., 2012. The Italian National Seismic Prevention Program. In: Proceedings of the 15th World Conference on Earthquake Engineering, Lisbon, Portugal.

- Douglas, J., 2010. Consistency of ground-motion predictions from the past four decades. *Bulletin of Earthquake Engineering* 8, 1515–1526.
- EFEHR, n.d. The European Facilities for Earthquake Hazard & Risk. www.efehr.org, [accessed 13 April, 2017].
- Erdik, M Durukal, E., 2008. Earthquake risk and its mitigation in Istanbul. *Natural Hazards* 44 (2), 181–197.
- Erdik, M., Aydinoglu, N., Fahjan, Y., Sesetyan, K., Demircioglu, M., Siyahi, B., Durukal, E., Ozbey, C., Biro, Y., Akman, H., Yuzugullu, O., 2003a. Earthquake risk assessment for Istanbul Metropolitan Area. *Earthquake Engineering and Engineering Vibration* 2 (1), 1–23.
- Erdik, M., Fahjan, Y., Orguz, O., AlcikH, Mert A, Gul M., 2003b. Istanbul Earthquake Rapid Response and the Early Warning System. *Bulletin of Earthquake Engineering* 1 (1), 157–163.
- Erduran, E., Lang, D., Lindholm, C., Toma-Danila, D., Balan, S., Ionescu, V., Aldea, A., Vacareanu, R., Neagu, C., 2012. Real-Time Earthquake Damage Assessment in the Romanian-Bulgarian Border Region. In: *Proceedings of the 15th World Conference on Earthquake Engineering*, Lisbon, Portugal.
- ERN-AL, 2009. Informe Técnico ERN-CAPRA-T1-3. Metodología de análisis probabilista del riesgo.
- ERSTA, 2010. Estudo do risco sísmico e de tsunamis do Algarve. Autoridade Nacional de Protecção Civil, Carnaxide, Portugal (in Portuguese).
- European Committee for Standardisation (CEN), 2005. EN 1998-1 Eurocode 8: Design of structures for earthquake resistance. Part 1: General rules, seismic actions and rules for buildings. European Committee for Standardization, Brussels, Belgium.
- FEMA, 2003. HAZUS-MH technical manual. Federal Emergency Management Agency, Washington, USA.
- Gamba, P., 2014. Global Exposure Database: scientific features. GEM Technical Report 2014-10. GEM Foundation, Pavia, Italy.
- Gardini, D., 1999. The Global Seismic Hazard Assessment Program (GSHAP) 1992-1999. *Annali di Geofisica* 42(6), :957–1230.
- Hancilar, U., Tuzun, C., Yenidogan, C. Erdik, M., 2010. ELER software—a new tool for urban earthquake loss assessment. *Natural Hazards and Earth Systems Sciences* 10, 2677–2696.
- Hoshiba, M., Kamigaichi, O., Saito, O., Tsukada, S., Hamada, N., 2008. Earthquake early warning starts nationwide in Japan. *EOS Transactions AGU* 89 (8), 73–74.
- Ioannou, I. , Rossetto, T., 2015. Empirical Fragility. In: Beer, M., Kouglioumtzoglou, I.A., Patelli, E., Siu-Kui Au, I. (Eds.), *Encyclopedia of Earthquake Engineering*. Springer, Berlin Heidelberg, pp. 976–986.
- Jaiswal, K., Aspinall, W., Perkins, D., Wald, D., Porter, K., 2012. Use of expert judgement elicitation to estimate seismic vulnerability of selected building types. In: *Proceedings of the 15th World Conference on Earthquake Engineering*, Lisbon, Portugal.
- Jaiswal, K., Wald, D., Porter, K., 2010. A global building inventory for earthquake loss assessment and risk management. *Earthquake Spectra* 26(3), 731–748.
- Kale, O., Akkar, S., 2013. A new perspective for selecting and ranking ground-motion prediction equations (GMPEs): the euclidian distance-based ranking method. *Bulletin of the Seismological Society of America*, 103(2A):1069–1084.
- Kale, O., Akkar, S., Ansari, A., Hamzehloo, H., 2015). A ground-motion predictive model for Iran and Turkey for horizontal PGA, PGV and 5 %-damped response spectrum: investigation of possible regional effects. *Bulletin of the Seismological Society of America* 105, 963–980.
- Khazai, B., Bendimerad, F., 2011. Risk and resiliency indicators, EMI Topical Report 565 TR-1 03.
- Martínez Solares, J., López Arroyo, A., 2004. The great historical 1755 earthquake. Effects and damage in Spain. *Journal of Seismology* 8, 275–294.
- Martins, L., Silva, V., Marques, M., Crowley, H., Delgado ,R. (2016). Development and assessment of damage-to-loss models for moment-frame reinforced concrete buildings. *Earthquake Engineering and Structural Dynamics* 45 (5),797–817.
- McGuire, R., (1976). Fortran program for seismic risk analysis. U.S. Geological Survey Open-File Report 76–67. <https://pubs.usgs.gov/of/1976/0067/report.pdf> [accessed 04 April, 2016]
- McGuire, R., 2004. Seismic hazard and risk analysis. Earthquake Engineering Research Institute, Oakland, USA.
- Mendes-Victor, L., Oliveira, C., Pais, I., Teves-Costa, P., 1994. Earthquake damage scenarios in Lisbon for disaster preparedness. In: Tucker, B.E., Erdik, M. Hwang, C.N. (Eds). *Issues in urban earthquake risk*. Kluwer Academic Publishers, Dordrecht, The Netherlands; Boston, USA, pp. 265–289.
- Molina, S., Lang, D., Lindholm, C., 2010. SELENA: An open-source tool for seismic risk and loss assessment using a logic tree computation procedure. *Computers & Geosciences* 36, 257–269.
- Mouroux, P., Le Brun, B., 2006. Presentation of RISK-UE Project. *Bulletin of Earthquake Engineering* 4, 323–339.
- NERA, n.d. NERA: Network of European Research Infrastructures for Earthquake Risk Assessment and Mitigation. <http://www.nera-eu.org/>, [accessed 12 April, 2017].
- NGDC, n.d. Natural Hazards Data, Images and Education. NOAA's National Centers for Environmental Information (NCEI) (formerly the National Geophysical Data Center). <https://www.ngdc.noaa.gov/hazard/hazards.shtml>, [accessed 12 April, 2017].
- Oliveira, C., Roca, A., Goula, X., 2006. Assessing and managing earthquake risk: geo-scientific and engineering knowledge for earthquake risk mitigation — development tools, techniques. Springer, Dordrecht, Netherlands.
- Pagani, M., Monelli, D., Weatherill, G., Danciu, L., Crowley, H., Silva, V., Henshaw, P., Butler, L., Nastasi, M., Panzeri, L., Simionato, M., Vigano, D., 2014. OpenQuake Engine: An open hazard (and risk) software for the Global Earthquake Model. *Seismological Research Letters* 85 (3),692–702.
- Papadopoulos, G.A., Arvanitides, A. 1996. Earthquake risk assessment in Greece. In: Schenk, V., (Ed.), *Earthquake hazard and risk, advances in natural & technological hazards research*, Kluwer Academic Publishers, USA.
- Pesaresi, M., Ehrlich, D., Ferri, S., Florczyk, A., Freire, S., Halkia, M., Julea, A., Kemper, T., Soille, P., Syrris, V., 2016. Operating procedure for the production of the Global Human Settlement Layer from Landsat data of the epochs 1975, 1990, 2000, and 2014. Technical Report EUR 27741 EN, Joint Research Center, Luxembourg.
- Petersen, M., Moschetti, M., Powers, P., Mueller, C., Haller, K., Frankel, A., Field, N., Chen, R., Rukstales, K.S., Luco, N., Wheeler, R., Williams, R., Olsen, A., 2015. The 2014 United States National Seismic Hazard Model. *Earthquake Spectra* 31(S1), S1–S30.
- Pitilakis, K., Crowley, H., Kaynia, A., 2014. SYNER-G: typology definition and fragility functions for physical elements at seismic risk: buildings, lifelines, transportation networks and critical facilities. Springer, Dordrecht, Netherlands.

- Pittore, M., Wieland, M., Fleming, K., 2016. Perspectives on global dynamic exposure modelling for geo-risk assessment. *Natural Hazards*, 86 (S1), 7-30.
- Porter, K., Farokhnia, K., Vamvatsioks, V., Cho, I., 2014. Guidelines for analytical vulnerability assessment of high-rise buildings. GEM Technical Report 2014. GEM Foundation, Pavia, Italy.
- REAKT, n.d. Strategies and tools for real time earthquake Risk Reduction. www.reaktproject.eu, [accessed 13 April, 2017].
- Renault, P., 2014. Approach and challenges for the seismic hazard assessment of nuclear power plants: the Swiss Experience. *Bollettino di Geoscienza Teorica ed Applicata* 55(1), 149-164.
- Rossetto, T., D'Ayala, D., Ioannou, I., Meslem, A., 2014b. Evaluation of existing fragility curves. In: Pitilakis, K., Crowley, H. and Kaynia, A.M. (eds), *SYNER-G: Typology definition and fragility functions for physical elements at seismic risk: buildings, lifelines, transportation networks and critical facilities*, Springer, Dordrech, Netherlands, pp. 47-93.
- Rossetto, T., Ioannou, I., Grant, D., 2014a. Guidelines for empirical vulnerability assessment. GEM Technical Report 2014-11. GEM Foundation, Pavia, Italy.
- Rossetto, T., Ioannou, I., Grant, D., 2015. Existing Empirical Fragility and Vulnerability Functions: Compendium and Guide for Selection (GEM Technical Report). GEM Foundation, Pavia, Italy.
- SAFER, n.d. Seismic early warning for Europe. <http://www.amrcenter.com/SAFER/index.htm>, [accessed 13 April, 2017].
- Scherbaum, F., Delavaud, E., Riggelsen, C., 2009. Model selection in seismic hazard analysis: an informationtheoretic perspective. *Bulletin of the Seismological Society of America* 99 (6), 3234-3247.
- Sengezer, B Koç, E., 2005. A critical analysis of earthquakes and urban planning in Turkey. *Disasters* 29 (2), 171-194.
- SHARE, n.d. Seismic Hazard Harmonization in Europe . www.share-eu.org, [accessed 13 April, 2017].
- SHEEC, n.d. The SHARE European Earthquake Catalogue. www.emidius.eu/SHEEC, [accessed 13 April, 2017].
- Silva, V., 2016. Critical Issues on Probabilistic Earthquake Loss Assessment. *Journal of Earthquake Engineering* 20 (8), 1322-1341.
- Silva, V., Crowley, H., Bazzurro, P., 2015. Exploring Risk-targeted Hazard Maps for Europe. *Earthquake Spectra* 32 (2), 1165-1186.
- Silva, V., Crowley, H., Pagani, M., Monelli D, Pinho, R., 2014b. Development of the OpenQuake engine, the Global Earthquake Model's open-source software for seismic risk assessment. *Natural Hazards* 72 (3), 1409-1427.
- Silva, V., Crowley, H., Pinho, R., Varum, H., 2014a. Seismic Risk Assessment for mainland Portugal. *Bulletin of Earthquake Engineering* 13 (2), 429-457.
- Spence, R., 2004. Risk and regulation: can improved government action reduce the impacts of natural disasters?. *Building Research & Information* 32 (5).
- Storchak, D., Di Giacomo, D., Endgdahl, E., Harris, J., Bondár, I., Lee, W., Bormann, P., Villaseñor, A., 2015. The ISC-GEM Global Instrumental Earthquake Catalogue (1900 — 2009): Introduction. *Physics of the Earth and Planetary Interiors* 239: 48-63.
- Strasser, F., Abrahamson, N., Bommer, J., 2009). *Sigma: Issues, Insights, and Challenges*. *Seismological Research Letters* 80, 41-56.
- STREST, n.d. Harmonized approach to stress tests for critical infrastructures against natural hazards. www.strest-eu.org, [accessed 13 April, 2017].
- Wald, D., Earle, P., Allen, T., Jaiswal, K., Porter, K., Hearne, M., 2012. Development of the US Geological Survey's PAGER system (prompt assessment of global earthquakes for response). In: *Proceedings of the 14th World Conference on Earthquake Engineering*, Lisbon, Portugal.
- Weatherill, G, Pagani, M., Garcia, J., 2016. Exploring Earthquake Databases for the Creation of Magnitude-Homogeneous Catalogues: Tools for Application on a Regional and Global Scale. *Geophysical Journal International* 206 (3), 165-276.
- Wieland, M., Pittore, M., Parolai, S. Zschau, J. (2012). Remote sensing and omnidirectional imaging for efficient building inventory data capturing: application within the Earthquake Model Central Asia. In: *Proceedings of the IEEE IGARSS 2012*, Munich, Germany.
- Woessner, J., Danciu, L., Giardini, D., Crowley H, Cotton F, Grunthal G, SHARE Consortium (2015) The 2013 European seismic hazard model: key components and results. *Bulletin of Earthquake Engineering* 13 (12), 3553-3596.
- Wu, Y.-M. Kanamori, H., 2008. Development of an Earthquake Early Warning System Using Real-Time Strong Motion Signals. *Sensors* 8 (1), 1-9.
- Yepes, C., Silva, V., Rossetto, T., D'Ayala, D., Ioannou, I., Meslem, A., Crowley, H., 2016. The Global Earthquake Model Physical Vulnerability Database. *Earthquake Spectra* 32 (4), 2567-2585.
- Zafarani, H., Mousavi, M., 2014. Applicability of different ground-motion prediction models for northern Iran. *Natural Hazards*, 73 (3), 1199-1228.
- Zollo, A., Iannaccone, G., Lancieri, M., Cantore, L., Convertito, V., Emolo, A., Festa, G., Galovic, F., Vassallo, M., Martino, C., Satriano, C., Gasparini, P., 2009. The earthquake early warning system in Southern Italy: methodologies and performance evaluation. *Geophysical Research Letters*, 36 (5).

3.2 Geophysical risk: volcanic activity

- Acocella, V., Di Lorenzo, R., Newhall, C., Scandone, R., 2015. An overview of recent (1988 to 2014) caldera unrest: Knowledge and perspectives. *Reviews of Geophysics* 53, 896-955.
- Aiuppa, A., Burton, M., Caltabiano, T., Giudice, G., Guerrieri, S., Liuzzo, M., Mure, F., Salerno, G., 2010. Unusually large magmatic CO₂ gas emissions prior to a basaltic paroxysm. *Geophysical Research Letters* 37, L17303.
- Aiuppa, A., Moretti, R., Federico, C., Giudice, G., Guerrieri, S., Liuzzo, M., Papale, P., Shinohara, H., Valenza, M., 2007. Forecasting Etna eruption by real time evaluation of volcanic gas composition. *Geology* 35, 1115-1118.
- Aiuppa, A., Tamburello, G., Di Napoli, R., Cardellini, C., Chiodini, G., Giudice, G., Grassa, F., Pedone, M., 2013. First observations of the fumarolic gas output from a restless caldera: Implications for the current period of unrest (2005-2013) at Campi Flegrei. *Geochemistry Geophysics Geosystems* 14, 4153-4169.
- Aspinall, W., 2010. A route to more tractable expert advice. *Nature* 463: 294-295.

- Aspinall, W.P., 2006. Structured elicitation of expert judgment for probabilistic hazard and risk assessment in volcanic eruptions. In: Mader, H.M., Coles, S. G., Connor, C.B., Connor, L.J. (Eds.), *Statistics in Volcanology*, Geological Society of London on behalf of IAVCEI, pp. 15–30.
- Auker, M., Sparks, R., Siebert, L., Croweller, H., Ewert, J., 2013. A statistical analysis of the global historical volcanic fatalities record. *Journal of Applied Volcanology* 2 (2), 1–24.
- Bagnardi, M., González, P.J., Hooper, A., 2016. High-resolution digital elevation model from tri-stereo Pleiades-1 satellite imagery for lava flow volume estimates at Fogo Volcano. *Geophysical Research Letters* 43, 6267–6275.
- Barclay, J., Haynes, K., Houghton, B., Johnston, D.M., 2015. Social processes and volcanic risk reduction. In: Sigurdsson, H., Houghton, B., McNutt, S., Rymer, H., Stix, J. (eds), *Encyclopedia of Volcanoes*. Academic Press, Amsterdam, Netherlands.
- Barclay, J., Haynes, K., Mitchell, T. O. M., Solana, C., Teeuw, R., Darnell, A., Croweller, H.S., Cole, P., Pyle, D., Lowe, C., Fearnley, C., 2014. Framing volcanic risk communication within disaster risk reduction: finding ways for the social and physical sciences to work together. In: Liverman, D., Pereira, C., Marker, B., (Eds.), *Communicating Environmental Geoscience*, Volume 305. Geological Society, London, UK, pp. 163–177.
- Bartolini, S., Bolós, X., Martí, J., Riera Pedra, E., Planaguma, L., 2015. Hazard assessment at the Quaternary La Garrotxa Volcanic Field (NE Iberia). *Natural Hazards* 78 (2), 1349–1367.
- Baxter, P. J., Aspinall, W.P., Neri, A., Zuccaro, G., Spence, R.J.S., Cioni, R., Woo, G., 2008. Emergency planning and mitigation at Vesuvius: A new evidence-based approach. *Journal of Volcanology and Geothermal Research* 178, 454–473.
- Baxter, P.J., Boyle, R., Cole, P., Neri, A., Spence, R., Zuccaro, G., 2005. The impacts of pyroclastic surges on buildings at the eruption of the Soufriere Hills volcano, Montserrat. *Bulletin of Volcanology* 67, 292–313.
- Bayarri, M.J., Berger, J.O., Calder, E.S., Patra, A.K., Pitman, E.B., Spiller, E.T., Wolpert, R.L., 2015. Probabilistic quantification of hazards: A methodology using small ensembles of physics based simulations and statistical surrogates. *International Journal for Uncertainty Quantification* 54, 297–325.
- Bebbington, M.S., Cronin S.J. 2010. Spatio-temporal hazard estimation in the Auckland volcanic field, New Zealand, with a new event-order model. *Bulletin of Volcanology*, 73, 55–72.
- Biass, S., Bonadonna, C., Di Traglia, F., Pistolesi, M., Rosi, M., Lestuzzi, P., 2016a. Probabilistic evaluation of the physical impact of future tephra fallout events for the Island of Vulcano, Italy. *Bulletin of Volcanology* 78, 37.
- Biass, S., Falcone, J.L., Bonadonna, C., Di Traglia, F., Pistolesi, M., Rosi, M., Lestuzzi, P., 2016b. Great Balls of Fire: A probabilistic approach to quantify the hazard related to ballistics — A case study at La Fossa volcano, Vulcano Island, Italy. *Journal of Volcanology and Geothermal Research* 325, 1–14.
- Biass, S., Scaini, C., Bonadonna, C., Folch, A., Smith, K., Höskuldsson, A., 2014. A multi-scale risk assessment for tephra fallout and airborne concentration from multiple Icelandic volcanoes - Part 1: Hazard assessment. *Natural Hazards and Earth System Sciences* 14, 2265–2287.
- Biggs, J., Anthony, E.Y., Ebinger, C.J. 2009. Multiple inflation and deflation events at Kenyan volcanoes, East African Rift. *Geology*, 37, 979–982.
- Biggs, J., Ebmeier, S.K., Aspinall, W.P., Lu, Z., Pritchard, M.E., Sparks, R.S.J., Mather, T.A., 2014. Global link between deformation and volcanic eruption quantified by satellite imagery. *Nature Communications* 5, 3471–3471.
- Bird, D.K., Gisladottir, G. and Dominey-Howes, D., 2010. Volcanic risk and tourism in southern Iceland: Implications for hazard, risk and emergency response education and training. *Journal of Volcanology and Geothermal Research*, 189, 33–48.
- Bird, P., 2003. An updated digital model of plate boundaries, *Geochemistry, Geophysics, Geosystems*, 4 (3), 1027.
- Birkmann, J., 2007. Risk and vulnerability indicators at different scales: applicability, usefulness and policy implications. *Environmental Hazards* 7, 20–31.
- Bonadonna C., Costa, A., 2012. Estimating the volume of tephra deposits: A new simple strategy, *Geology*, 40 (5), 415–418.
- Bonadonna, C., 2006. Probabilistic modelling of tephra dispersal. In: Mader, H., Cole, S., Connor C.B. (Eds.), *Statistics in volcanology*. IAVCEI Series Volume 1. Geological Society, London, UK, pp. 243–259.
- Bonadonna, C., Biass, S., Costa, A., 2015. Physical characterization of explosive volcanic eruptions based on tephra deposits: Propagation of uncertainties and sensitivity analysis. *Journal of Volcanology and Geothermal Research* 296, 80–100.
- Bonadonna, C., Folch, A., Loughlin, S. and Puempel, H., 2012. Future developments in modelling and monitoring of volcanic ash clouds: Outcomes from the first IAVCEI-WMO workshop on Ash Dispersal Forecast and Civil Aviation. *Bulletin of Volcanology* 74, 1–10.
- Boué, A., Lesage, P., Cortés, G., Valette, B., Reyes Dávila, G., Arámbula Mendoza, R., Budi Santoso, A., 2016. *Journal of Volcanology and Geothermal Research* 327, 622–633.
- Bretton R.J., Gottsman J.H., Aspinall W.P., Christie R.B., 2015. Implications of legal scrutiny processes (including the L'Aquila trial and other recent court cases) for future volcanic risk governance. *Journal of Applied Volcanology* 4 (1), 18.
- Brown, S.K., Croweller, H.S., Sparks, R.S.J., Cottrell E., Deligne, N.I., Ortiz Guerrero, N., Hobbs, L., Kiyosugi, K., Loughlin, S.C., Siebert, L., Takarada, S., 2014. Characterisation of the Quaternary eruption record: analysis of the Large Magnitude Explosive Volcanic Eruptions (LaMEVE) database. *Journal of Applied Volcanology* 3 (5), 22.
- Brown, S.K., Loughlin, S.C., Sparks, R.S.J., Vye-Brown, C., Barclay, J., Calder, E., Cottrell, E., Jolly, G., Komorowski, J.C., Mandeville, C., Newhall, C., Palma, J., Potter, S., Valentine, G., 2015. Global volcanic hazard and risk. In: Loughlin, S.C., Sparks, R.S.J., Brown, S.K., Jenkins, S.F., Vye-Brown, C., (eds), *Global Volcanic Hazards and Risk*. Cambridge University Press, Cambridge, pp. 81 – 173.
- Calder, E.S., Wagner, K. Ogburn, S.E. (2015). Volcanic hazard maps. In: Loughlin, S.C., Sparks, R.S.J., Brown, S.K., Jenkins, S.F., Vye-Brown, C. (eds), *Global Volcanic Hazards and Risk*. Cambridge University Press, Cambridge, UK.
- Cannavò, F., Camacho, A. G., González, P. J., Mattia, M., Puglisi, G. and Fernández, J., 2015. Real Time Tracking of Magmatic Intrusions by means of Ground Deformation Modeling during Volcanic Crises, *Scientific Reports* 5.
- Carlsen, H. K., Hauksdottir, A., Valdmarisdottir, U. A., Gíslason, T., Einarsdottir, G., Runolfsson, H., Briem, H., Finnbjornsdottir, R. G., Gudmundsson, S. and Kolbeinsson, T. B., 2012. Health effects following the Eyjafjallajökull volcanic eruption: a cohort study. *BMJ Open*, 2, e001851.
- Cashman, K.V., Stephen, R., Sparks, J., 2013. How volcanoes work: A 25 year perspective. *Bulletin of the Geological Society of*

- America 125, 664-690.
- Charbonnier S.J., Germa A., Connor C.B., Gertisser R., Preece K., Komorowski J.C., Lavigne F., Dixon T. and Connor, L. undefined, 2013. Evaluation of the impact of the 2010 pyroclastic density currents at Merapi volcano from high-resolution satellite imagery, field investigations and numerical simulations. *Journal of Volcanology and Geothermal Research* 261 (295), 315.
- Chiodini, G., Caliro, S., De Martino, P., Avino, R., Gherardi, F., 2012. Early signals of new volcanic unrest at Campi Flegrei caldera? Insights from geochemical data and physical simulations. *Geology* 40, 943-946.
- Chiodini, G., Paonita, A., Aiuppa, A., Costa, A., Caliro S., De Martino, P., Acocella, V., Vandemeulebrouck, J., 2016. Magmas near the critical degassing pressure drive volcanic unrest towards a critical state. *Nature Communications* 7, 13712.
- Chiodini, G., Pappalardo, L., Aiuppa, A., and Caliro, S., 2015. The geological CO₂ degassing history of a long-lived caldera. *Geology* 43 (9), 767-770.
- Conde, V., Robidoux, P., Avar, G., Galle, B., Aiuppa, A., Muñóz, A., 2013. Measurements of SO₂ and CO₂ by combining DOAS, Multi-GAS and FTIR: study cases from Turrialba and Telica volcanoes. *International Journal of Earth Sciences* 103 (8), 2335-2347.
- Connor, C., Bebbington, M., Marzocchi, W., 2015. Probabilistic Volcanic Hazard Assessment. In: Sigurdsson, H., Houghton, B., McNutt, S., Rymer, H., Stix, S., (eds.), *Encyclopedia of Volcanoes*. Academic Press, Amsterdam, Netherlands, pp. 897-910.
- Connor, L. J., Connor, C. B., Meliksetian, K., Savov, I. 2012. Probabilistic approach to modelling lava flow inundation: a lava flow hazard assessment for a nuclear facility in Armenia. *Journal of Applied Volcanology*, 1, 3.
- Coppola, D., Laiolo, M., Cigolini, C., Delle Donne, D., Ripepe, M., 2015. Enhanced volcanic hot-spot detection using MODIS IR data: results from the MIROVA system. In: Harris, A.J.L., De Groeve, T., Garel, F., Carn, S.A. (Eds.), *Detecting, modelling and responding to effusive eruptions*. Geological Society, London, UK.
- Cottrell, E., 2014. Global Distribution of Active Volcanoes. In: Papale, P. (ed.) *Volcanic Hazards, Risks and Disasters*. Academic Press, pp. 1-18.
- Craig, H., Wilson, T., Stewart, C., Outes, V., Villarosa, G., Baxter, P., 2016. Impacts to agriculture and critical infrastructure in Argentina after ashfall from the 2011 eruption of the Cordón Caulle volcanic complex: an assessment of published damage and function thresholds. *Journal of Applied Volcanology* 5, 7.
- Cutter, S.L., 2013. Vulnerability. In: *Encyclopedia of Natural Hazards, Encyclopedia of Earth Sciences Series*. Springer-Verlag, Berlin, Heidelberg, Germany.
- De Natale, G., Troise, C., Pingue, F., Mastrolorenzo, G., Pappalardo, L., Battaglia, M., Boschi, E., 2006. The Campi Flegrei caldera: unrest mechanisms and hazards. In: Troise, C., De Natale, G., Kilburn, C.R.J. (eds.), *Mechanisms of Activity and Unrest at Large Calderas*. Geological Society, London, UK, pp. 25-45.
- Deligne, N.I., Coles, S.G., Sparks, R.S.J., 2010. Recurrence rates of large explosive volcanic eruptions. *Journal of Geophysical Research: Solid Earth* 115, B06203.
- Doyle, E. H., McClure, J., Johnston, D. M., Paton, D., 2014. Communicating likelihoods and probabilities in forecasts of volcanic eruptions. *Journal of Volcanology and Geothermal Research* 272, 1-15.
- Doyle, E. H., Paton, D. Johnston, D. M., 2015. Enhancing scientific response in a crisis: evidence-based approaches from emergency management in New Zealand. *Journal of Applied Volcanology* 4,1.
- Druitt, T. H., Edwards, L., Mellors, R. M., Pyle, D. M., Sparks, R. S. J., Lanphere, M., Davies, M., Barreiro, B., 1999. Santorini Volcano. Geological Society, London, UK.
- Elissondo, M., Baumann, V., Bonadonna, C., Pistolesi, M., Cioni, R., Bertagnini, A., Biass, S., Herrero, J-C., Gonzalez, R., 2016. Chronology and impact of the 2011 Cordón Caulle eruption, Chile. *Natural Hazards and Earth Systems Science* 16, 675-704.
- Engwell, S. L., Sparks, R. S. J., Aspinall, W. P., 2013. Quantifying uncertainties in the measurement of tephra fall thickness. *Journal of Applied Volcanology* 2 (1), 1-12.
- Falsaperla, S. and Neri, M., 2015. Seismic footprints of shallow dyke propagation at Etna, Italy, *Scientific Reports*, 5, 11908.
- Fearnley, C.J., 2013. Assigning a volcano alert level: negotiating uncertainty, risk, and complexity in decision-making processes. *Environment and Planning A*, 45, 1891-1911.
- Fee, D., Matoza, R.S., 2013. An overview of volcano infrasound: From Hawaiian to plinian, local to global. *Journal of Volcanology and Geothermal Research* 249, 123-130.
- Flower, V.J.B., Oommen, T., Carn, S.A., 2016. Improving global detection of volcanic eruptions using the Ozone Monitoring Instrument (OMI), *Atmospheric Measurement Techniques* 9, 5487-5498.
- Frey Mueller, J. T., Murray, J. B., Rymer, H., Locke, C. A., 2015, Ground deformation, Gravity and Magnetics, 1101-1123 In: Sigurdsson, H., Houghton, B., McNutt, S., Rymer, H. and Stix, S., (eds.), *Encyclopedia of Volcanoes*. Academic Press, USA.
- Gaillard, J. C., 2008. Alternative paradigms of volcanic risk perception: The case of Mt. Pinatubo in the Philippines. *Journal of Volcanology and Geothermal Research* 172, 315-328.
- Galderisi, A., Bonadonna, C., Delmonaco, G., Ferrara, F. F., Menoni, S., Ceudech, A., Biass, S., Frischknecht, C., Manzella, I., Minucci, G., Gregg, C., 2011. Vulnerability assessment and risk mitigation: the case of Vulcano Island, Italy. In: *Proceedings of the Second World Landslide Forum*, Rome.
- Gardner, C.A., Guffanti, M.C., 2006. U.S. Geological Survey's alert notification system for volcanic activity. *U.S. Geological Survey Fact Sheet* 2006-3139.
- Gislason, S. R., Stefansdóttir, G., Pfeffer, M. A., Barsotti, S., Johansson, Th, Galecka, I., Bali, E., Sigmarsson, O., Stefansson, A., Keller, N. S., Sigurdsson, A., Bergsson, B., Galle, B., Jacobo, V. C., Arellano, S., Aiuppa, A., Jonasdóttir, E. B., Eiríksdóttir, E. S., Jakobson, S., Gudfinnson, G. H., Halldorsson, S. A., Gunnarson, H., Haddadi, B., Jonsdóttir, I., Thordarson, T., Riishus, M., Hognadóttir, T., Durig, T., Pedersen, G. B.M., Hoskuldsson, A., Gudmundsson, M.T., 2015. Environmental pressure from the 2014-15 eruption of Bardarbunga volcano, Iceland. *Geochemical Perspectives Letters* 1.
- Gudmundsson, M.T., Jónsdóttir, K., Hooper, A., Holohan, E.P., Halldórsson, S.A., Ófeigsson, B.G., Cesca, S., Vogfjörð, K.S., Sigmundsson, F., Högnadóttir, T., Einarsson, P., Sigmarsson, O., Jarosch, A.H., Jónasson, K., Magnússon, E., Hreinsdóttir, S., Bagnardi, M., Parks, M.M., Hjörleifsdóttir, V., Pálsson, F., Walter, T.R., Schöpfer, M.P.J., Heimann S., Reynolds, H.I., Dumont, S., Bali, E., Gudfinnsson, G.H., Dahm, T., Roberts, M.J., Hensch, M., Belart, J.M.C., Spaans, K., Jakobsson, S., Gudmundsson, G.B., Fridriksdóttir, H.M., Drouin, V.,

- Dürrig, T., Aðalgeirsdóttir, G., Riishuus, M.S., Pedersen, G.B.M., Van Boeckel, T., Oddsson, B., Pfeffer, M.A., Barsotti, S., Bergsson, B., Donovan, A., Burton, M.R., Aiuppa, A., 2016. Gradual caldera collapse at Bárðarbunga volcano, Iceland, regulated by lateral magma outflow. *Science* 353(6296), aaf8988.
- Gudmundsson, M.T., Thordarson, T., Höskuldsson, A., Larsen, G., Björnsson, H., Prata, F., Oddson, B., Magnusson, E., Hognadóttir, T., Petersen, G., Hayward, C. L., Stevenson, J. A., Jonsdóttir, I., 2012. Ash generation and distribution from April-May 2010 eruption of Eyjafjallajökull, Iceland. *Scientific Reports* 2 (572), 1-12.
- Guffanti, M., Casadevall, T.J., Budding, K., 2010. Encounters of aircraft with volcanic ash clouds: a compilation of known incidents, 1953-2009. *US Geological Survey Data Series* 545, 12- 12.
- Harris, A., Dehn, J., Patrick, M., Calvari, S., Ripepe, M., and Lodato, L., 2005. Lava effusion rates from hand-held thermal infrared imagery; an example from the June 2003 effusive activity at Stromboli. *Bulletin of Volcanology*, 68 (2), 107-117.
- Harris, A.J.L., De Groeve, T., Garel, F. Carn, S. E. (eds), 2016. *Detecting, Modelling and Responding to Effusive Eruptions*. Geological Society, London, UK.
- Hartley, M. E., Morgan, D. J., MacLennan, J., Edmonds, M., Thordarson, T., 2016. Tracking timescales of short-term precursors to large basaltic fissure eruptions through Fe-Mg diffusion in olivine. *Earth and Planetary Science Letters* 439, 58-70.
- Hasegawa, Y., Sugai, A., Hayashi, Yo., Hayashi, Yu., Saito, S., Shimbori, T., 2015. Improvements of volcanic ash fall forecasts issued by the Japan Meteorological Agency. *Journal of Applied Volcanology Society and Volcanoes* 4(2).
- Hayes, J., Wilson, T.M., Deligne, N.I., Cole, J., Hughes, M., 2017. A model to assess tephra clean-up requirements in urban environments. *Journal of Applied Volcanology* 6,1.
- Haynes, K., Barclay, J., Pidgeon, N., 2008a. The issue of trust and its influence on risk communication during a volcanic crisis. *Bulletin of Volcanology* 70, 605-621.
- Haynes, K., Barclay, J., Pidgeon, N., 2008b. Whose reality counts? Factors affecting the perception of volcanic risk. *Journal of Volcanology and Geothermal Research* 172, 259- 272.
- Haynes, K., Barclay, J., Pidgeon, N., 2007. Volcanic hazard communication using maps: an evaluation of their effectiveness. *Bulletin of Volcanology* 70, 123-138.
- Hernández, P.A., Pérez, N.M., Varekamp, J.C., Henriquez, B., Hernández, A., Barrancos, J., Padrón, E., Calvo, D., Melián, G., 2007. Crater Lake Temperature Changes of the 2005 Eruption of Santa Ana Volcano, El Salvador, Central America. *Pure and Applied Geophysics* 164, 2507-2522.
- Hicks, A. and Few, R., 2015. Trajectories of social vulnerability during the Soufriere Hills Volcanic crisis, *Journal of Applied Volcanology* 4, 10.
- Hicks, A., Barclay, J., Mark, D. F., Loughlin, S., 2012. Tristan da Cunha: Constraining eruptive behaviour using the ⁴⁰Ar/³⁹Ar dating technique. *Geology* 40, 723-726.
- Hicks, A., Barclay, J., Simmons, P., Loughlin, S., 2014. An interdisciplinary approach to volcanic risk reduction under conditions of uncertainty: a case study of Tristan da Cunha. *Natural Hazards and Earth System Sciences* 14, 1871-1887.
- Hincks, T. K., Komorowski, J.-C., Sparks, S. R., Aspinall, W. P. 2014. Retrospective analysis of uncertain eruption precursors at La Soufrière volcano, Guadeloupe, 1975-77: volcanic hazard assessment using a Bayesian Belief Network approach. *Journal of Applied Volcanology* 3, 1-26.
- Horwell, C. and Baxter, P. J., 2006. The respiratory health hazards of volcanic ash: a review for volcanic risk mitigation. *Bulletin of Volcanology* 69, 1-24.
- IAVCEI Subcommittee for Crisis Protocols, 1999. Professional conduct of scientists during volcanic crises. *Bulletin of Volcanology* 60, 323-334.
- Ilyinskaya, E., Larsen, G., Gudmundsson, M.T., 2015. The Catalogue of Icelandic Eruptions.
- IMO, 2016. Monitoring of Hekla. Icelandic met office. <http://en.vedur.is/about-imo/news/monitoring-hekla>, [accessed 13 April, 2017].
- IMO, n.d. Volcanic gas detected. Icelandic met office <http://en.vedur.is/pollution-and-radiation/volcanic-gas/>, [accessed 13 April, 2017].
- INGV, n.d. INGV Sezione di Catania. Osservatorio Etneo. <http://www.ct.ingv.it/it/simulazione-dispersione-ceneri.html>, [accessed 13 April, 2017].
- Integrated Research on Disaster Risk, 2011. *Forensic Investigations of Disasters: The FORIN Project*. Integrated Research on Disaster Risk Programme, Beijing, China.
- Jenkins, S. F., Phillips, J. C., Price, R., Feloy, K., Baxter, P. J., Hadmoko D. S., de Bézilal, E., 2015b. Developing building-damage scales for lahars: application to Merapi volcano, Indonesia. *Bulletin of Volcanology* 77, 75.
- Jenkins, S., Magill, C., McAneney, J., Blong, R., 2012. Regional ash fall hazard I: a probabilistic assessment methodology. *Bulletin of Volcanology* 74, 1699-1712.
- Jenkins, S.F., Spence, R.J.S., Fonseca, J.F.B. D., Solidum, R.U., Wilson, T.M., 2014. Volcanic risk assessment: Quantifying physical vulnerability in the built environment. *Journal of Volcanology and Geothermal Research* 276, 105-20
- Jenkins, S.F., Wilson, T., Magill, C., Miller, V., Stewart, C., Blong, R., Marzocchi, W., Boulton, M., Bonadonna, C. and Costa, A., 2015a. Volcanic ash fall hazard and risk. In: Loughlin, S.C., Sparks, R.S.J., Brown, S.K, Jenkins, S.F., Vye-Brown, C. (Eds), *Global Volcanic Hazards and Risk*. Cambridge University Press, Cambridge, UK.
- Johnson, J. B. and Ripepe, M., 2011. Volcano infrasound: A review. *Journal of Volcanology and Geothermal Research* 206, 61-69.
- Jones, R., Manville, V., Andrade D., 2015 Probabilistic analysis of rainfall triggered lahar initiation at Tungurahua Volcano. *Bulletin of Volcanology* 77, 68.
- Kristmansdóttir, H., Björnsson, A., Pálsson, S., Sveinbjörnsdóttir, A.E., 1999. The impact of the 1996 subglacial eruption in Vatnajökull on the river Jökulsá á Fjöllum, North Iceland. *Journal of Volcanology and Geothermal Research* 92(3-4), 359-372.
- Lamb, O.D., De Angelis, S., Lavalée, Y., 2015. Using infrasound to constrain ash plume rise. *Journal of Applied Volcanology* 4, 20.
- Lechner, P., Tupper, A., Guffanti, M., Loughlin, S., Casadevall, T., 2017. Volcanic ash and aviation — the challenges of real-time, global communication of a natural hazard. *Advances in Volcanology*. Springer Berlin Heidelberg, 1-14
- Lindsay, J., Marzocchi, W., Jolly, G., Constantinescu, R., Selva, J., Sandri, L., 2010. Towards real-time eruption forecasting in the

- Auckland Volcanic Field: application of BET_EF during the New Zealand National Disaster Exercise 'Rauumoko'. *Bulletin of Volcanology* 72, 185.
- Lindsay, J.M, Marzocchi, W., Jolly, G, Constantinescu, R., Selva, J., Sandri, L., 2010. Towards real-time eruption forecasting in the Auckland volcanic field: application of BET_EF during the New Zealand national disaster Exercise 'Rauumoko'. *Bulletin of Volcanology* 2, 185-204.
- Loughlin, S., Baxter, P., Aspinall, W., Darroux, B., Harford, C. and Miller, A., 2002. Eyewitness accounts of the 25 June 1997 pyroclastic flows and surges at Soufrière Hills Volcano, Montserrat, and implications for disaster mitigation. In: Druitt, T.H., and Kokelaar, B.P. (eds.), *The Eruption of Soufriere Hills Volcano, Montserrat, from 1995 to 1999*. Geological Society, London, UK, *Memoirs* 21, 211-230.
- Macedonio, A., Costa, A. Folch, A., 2008. Ash fallout scenarios at Vesuvius: Numerical simulations and implications for hazard assessment. *Journal of Volcanology and Geothermal Research* 178, 366-377.
- Magill, C., Wilson, T., Okada, T., (2013). Observations of tephra fall impacts from the 2011 Shinmoedake eruption, Japan. *Earth, Planets and Space* 65, 677-698.
- Marchetti, E., Innocenti, L., Olivieri, G., Lacanna, G., Ripepe, M., 2016. Long-range infrasound monitoring of eruptive volcanoes. *Geophysical Research Abstracts*, 18, EGU2016-13157. EGU General Assembly 2016.
- Martí, J., Aspinall, W. P., Sobradelo, R., Felpeto, A., Geyer, A., Ortiz, R., Baxter, P., Cole, P., Pacheco, J., Blanco, M. J., Lopez, C., 2008. A long-term volcanic hazard event tree for Teide-Pico Viejo stratovolcanoes (Tenerife, Canary Islands). *Journal of Volcanology and Geothermal Research* 178, 543-552.
- Marzano, F. S., Picciotti, A., Di Fabio, S., Montopoli, M., Mereu, L., Degruyter, W., Bonadonna, C. and Ripepe, M., 2016. Near-Real-Time Detection of Tephra Eruption Onset and Mass Flow Rate Using Microwave Weather Radar and Infrasonic Arrays. *IEEE Transactions on Geoscience and Remote Sensing*.
- Marzano, F., Picciotti, E., Montopoli, M. Vulpiani, G., 2013. Inside Volcanic Clouds: Remote Sensing of Ash Plumes Using Microwave Weather Radars. *Bulletin of the American Meteorological Society* 94, 1567-1586.
- Marzocchi W., Newhall C., Woo, G., 2012. The Scientific Management of Volcanic Crises. *Journal of Volcanology & Geothermal Research* 247-248, 181-189.
- Marzocchi, W, Sandri L, Selva, J., 2008. BET_EF: a probabilistic tool for long- and short-term eruption forecasting. *Bulletin of Volcanology* 70, 623-632.
- Marzocchi, W. and Bebbington, M. S., 2012. Probabilistic eruption forecasting at short and long time scales. *Bulletin of Volcanology* 74, 1777-1805.
- Marzocchi, W., Neri, A., Newhall, C. G. and Papale, P., 2007. Probabilistic volcanic hazard and risk assessment, *Eos Transactions, AGU*, 88 (32), 318.
- McNutt, S. Thompson, G., Johnson, J. and Fee, D., 2015. Seismic and infrasonic monitoring. *Encyclopedia of Volcanoes*, Edition 2nd, Chapter 63, Elsevier, pp.1071-1099.
- Menoni, S., Molinari, D., Parker, D., Ballio, F., Tapsell, S., 2012. Assessing multifaceted vulnerability and resilience in order to design risk-mitigation strategies. *Natural Hazards* 64, 2057-2082.
- Mothes, P.A., Yepes, H. A., Hall, M. L., Ramon, P.A., Steele, A. L. and Ruiz, M.C., 2015. The scientific-community interface over the fifteen-year eruptive episode of Tungurahua Volcano, Ecuador. *Journal of Applied Volcanology* 4,9.
- Nadeau, P.A., Palma, J.L., Waite, G.P., 2011. Linking volcanic tremor, degassing and eruption dynamics via SO₂ imaging. *Geophysical Research Letters* 38, L01304.
- Newhall, C. and Hoblitt, R., 2002. Constructing event trees for volcanic crises. *Bulletin of Volcanology* 64, 3-20.
- Newhall, C. G., Self, S., 1982. The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism. *Journal of Geophysical Research* 87, 1231-1231.
- Newhall, C.G., Hendley II, J.W., Stauffer, P.H., 1997. Benefits of volcano monitoring far outweigh costs — the case of Mount Pinatubo, *Fact Sheet* 115-97.
- Ogburn, S.E., Loughlin, S.C., Calder, E. S., 2015. The association of lava dome growth with major explosive activity (VEI>4): Dome-Haz, a global dataset. *Bulletin of Volcanology* 77, 40.
- Oppenheimer, C.M.M., Fischer, T. and Scaillet, B., 2013. Volcanic degassing: process and impact. *Treatise on Geochemistry* 4, 111-179.
- Orsi, G., Di Vito, M., Isaia R., 2004. Volcanic hazard assessment at the restless Campi Flegrei caldera. *Bulletin of Volcanology* 66, 514-530.
- Pallister, J. and McNutt, S. R., 2015. Synthesis of Volcano Monitoring, In: Sigurdsson, H., Houghton, B., McNutt, S., Rymer, H. And Stix, J. (Eds), *The Encyclopedia of Volcanoes*, Elsevier, 1424pp.
- Pallister, J. and Surono, 2015. Forecasting the November 2010 eruption of Merapi, Indonesia. In: Loughlin, S.C., Sparks, R.S.J., Brown, S.K., Jenkins, S.F., Vye-Brown, C. (eds), *Global Volcanic Hazards and Risk*. Cambridge University Press, Cambridge, UK.
- Pankhurst, M. J., Dobson, K. J., Morgan, D. J., Loughlin, S. C., Thordarson, Th., Lee, P. D. and Courtois, L., 2014. Monitoring the magmas fuelling volcanic eruption in near-real-time using X-ray micro-computed tomography. *Journal of Petrology* 55 (3), 671-684.
- Papadopoulos, G.A., Orfanogiannaki, K., 2005. Long-term prediction of the next eruption in Thera volcano from conditional probability estimates. In: Fytikas M., Vougioukalakis, G.E., (eds.), *The South Aegean Volcanic Arc*. Elsevier, pp. 211-216.
- Parks, M.M., Moore, J. D.P., Papanikolaou, X., Biggs, J., Mather, T.A., Pyle, D. M., Raptakis, C., Paradissis, D., Hooper, A., Parsons, B., Nomikou, P., 2015. From quiescence to unrest: 20 years of satellite geodetic measurements at Santorini volcano, Greece, *Journal of Geothermal Research B: Solid Earth* 120(2), 1309-1328.
- Phillipson, G., Sobradelo, R., Gottsmann, J., 2013. Global volcanic unrest in the 21st century: an analysis of the first decade. *Journal of Volcanology and Geothermal Research* 264, 183- 196.
- Potter, S. H., Jolly, G. E., Neall, V. E., Johnston, D. M., Scott, B. J., 2014. Communicating the status of volcanic activity: revising New Zealand's volcanic alert level system. *Journal of Applied Volcanology* 3, 1-16.
- Pyle, D. 2015. Sizes of Volcanic Eruptions. In: Sigurdsson, H., Houghton, B., McNutt, S., Rymer, H. and Stix, J. *The Encyclopedia of Volcanoes*, 2nd Edition, 257-264.

- Ragona, M., Hannstein, F. and Mazzocchi, M., 2011. The impact of volcanic ash crisis on the European Airline industry. In: Alemanno, A., (ed.), *Governing Disasters: The Challenges of Emergency Risk regulations*. Edward Elgar Publishing, Cheltenham, UK/Northampton, USA.
- Ricci T., Nave R., Barberi F., 2013. Vesuvio civil protection exercise MESIMEX: survey on volcanic risk perception. *Annals of Geophysics* 56(4), S0452.
- Richter, N., Favalli, M., de Zeeuw-van Dalfsen, E., Fornaciai, A., Fernandes, R., Pérez, N. M., Levy, J., Victória, S. S., Walter, T. R., 2016. Lava flow hazard at Fogo Volcano, Cabo Verde, before and after the 2014-15 eruption. *Natural Hazards and Earth System Science* 16, 1925-1951.
- Ripepe, M., Bonadonna, C., Folch, A., Delle Donne, D., Lacanna, G., Marchetti, E., Höskuldsson, A., 2013. Ash-plume dynamics and eruption source parameters by infrasound and thermal imagery: the 2010 Eyjafjallajökull eruption. *Earth and Planetary Science Letters* 366, 112-121.
- Roberts, M. J., Linde, A. T., Vogfjord, K. S., Sacks, S., 2011. Forecasting Eruptions of Hekla Volcano, Iceland, using Borehole Strain Observations. *Geophysical Research Abstracts* 13, EGU2011-14208.
- Saltogiani, V., Stiros, S. C., Newman, A. V., Flanagan, K., Moschas, F., 2014. Time-space modeling of the dynamics of Santorini volcano (Greece) during the 2011-2012 unrest. *Journal of Geophysical Research: Solid Earth* 119, 8517-8537
- Savov, I., Meliksetian, K., Connor, C., Karakhanian, A., Sugden, P., Navasardyan, G., Halama, R., Ishizuka, O., Connor, L., Karapetian, S., 2016. Evolution of Pleistocene to Holocene eruptions in the Lesser Caucasus Mts: Insights from geology, petrology, geochemistry and geochronology. *Geophysical Research Abstracts* 18, EGU2016-6377.
- Scandone, R., Arganese, G., Galdi, F., 1993. The evaluation of volcanic risk in the Vesuvian area. *Journal of Volcanology and Geothermal Research* 58, 263-271.
- Schmidt, A., Leadbetter, S., Theys, N., Carboni, E., Witham, C. S., Stevenson, J. A., Birch, C. E., Thordarson, Th., Turnock, S., Barsotti, S., Delaney, L., eng, W., Grainger, R. G., Hort, M. C., Hoskuldsson, A., Ialongo, I., Ilyinskaya, E., Johansson, Th., Kenny, P., Mather, T. A., Richards, N. A. D., Shepherd, J., 2015. Satellite detection, long-range transport, and air quality impacts of volcanic sulphur dioxide from the 2014-2015 flood lava eruption at Bardarbunga (Iceland). *Journal of Geothermal Research: Atmospheres*,
- Schmidt, A., Ostro, B., Carslaw, K. S., Wilson, M., Thordarson, T., Mann, G. W. and Simmons, A. J., 2011. Excess mortality in Europe following a future Laki-style Icelandic eruption. *Proceedings of the National Academy of Sciences* 108, 15710-15715.
- Scientific advisory board of the Icelandic civil protection, 2015. Volcanic activity in the Bardarbunga system. http://en.vedur.is/media/jar/Factsheet_Bardarbunga_20150127.pdf, [accessed 13 April, 2017].
- Segall, P. 2013. Volcano deformation and eruption forecasting. *Geological Society, Special Publications* 380, 85-106.
- Selva, J., Costa, A., Marzocchi, W., Sandri, L., 2010. BET_VH: Exploring the influence of natural uncertainties on long-term hazard from tephra fallout at Campi Flegrei (Italy). *Bulletin of Volcanology* 72, 717.
- Selva, J., Marzocchi, W., Sandri, L., Costa, A. 2015. Operational Short-term Volcanic Hazard Analysis: Methods and Perspectives. In: Papale (Ed), *Volcanic Hazards, Risks and Disasters*, 233-260.
- Selva, J., Orsi, G., Di Vito, M. A., Marzocchi, W., Sandri, L., 2012. Probability hazard map for future vent opening at the Campi Flegrei caldera, Italy. *Bulletin of Volcanology*, 74, 497-510.
- Siebert, L., Cottrell, E., Venzke, E., Andrews, B., 2015. Earth's volcanoes and Their Eruptions: An Overview. In: Sigurdsson, H., Houghton, B., Rymer, H., Stix, J., McNutt, (Eds.). *Encyclopedia of Volcanoes*, Academic Press.
- Siebert, L., Simkin, T. and Kimberley, P., 2010. *Volcanoes of the World*, 3rd edn, University of California Press, Berkeley, USA.
- Sigmundsson, F., Hooper, A., Hreinsdottir, S., Vojfjord, K.S., Ofeigsson, B.G., Heimisson, E.R., Dumont, S., Parks, M., Spaans, K., Gudmundsson, G.B., Drouin, V., Arnadottir, T., Jonsdottir, K., Gudmundson, M.T., Hognadottir, T., Fridriksdottir, H.M., Hensch, M., Einarsson, P., Magnússon, E., Samsonov, S., Brandsdottir, B., White, R.S., Agustsdottir, T., Greenfield, T., Green, R.G., 2015. Segmented lateral dyke growth in a rifting event at Bardarbunga volcanic system, Iceland. *Nature* 517, 191-195.
- Sigmundsson, F., Hreinsdottir, A., Hooper, A., Arnadottir, T., Pedersen, R., Roberts, M. J., Oskarsson, N., Auriac, A., Decriem, J., Einarsson, P., Geirsson, H., Hensch, M., Ofeigsson, B. G., Sturkell, E., Sveinbjornsson, H., Feigi, K. L., 2010. Intrusion triggering of the 2010 Eyjafjallajökull explosive eruption. *Nature* 468, 426-430.
- Silva, C., Viveiros, F., Ferreira, T., Gaspar, J.L., Allard, P., 2015. Diffuse soil emanations of radon and hazard implications at Furnas Volcano, São Miguel Island (Azores). *Geological Society London, Memoirs* 44, 197-211.
- Smithsonian Institution, 2013. *Volcanoes of the World 4.0*. <http://www.volcano.si.edu> [accessed 06 April, 2016]
- Solana, M. C., Kilburn, C.R.J. and Rolandi, G., 2008. Communicating eruption and hazard forecasts on Vesuvius, Southern Italy. *Journal of Volcanology and Geothermal Research* 172, 308- 314.
- Spaans, K., Hooper, A., 2016. InSAR processing for volcano monitoring and other near-real time applications, *Journal of Geophysical Research Solid Earth* 121 (4), 2947-2960.
- Sparks, R.S.J., 2003. Forecasting volcanic eruptions. *Earth and Planetary Science Letters* 210, 1-15.
- Sparks, R.S.J., Aspinall, W.P., Crossweller, H.S. and Hincks, T.K., 2013. Risk and uncertainty assessment of volcanic hazards. In: Rougier, J., Sparks, R. S. J., Hill, L., (Eds.). *Risk and Uncertainty Assessment for Natural Hazards*. Cambridge University Press, Cambridge, UK, 364-397.
- Sparks, R.S.J., Biggs, J. and Neuberg, J., 2012. Monitoring volcanoes. *Science*, 335, 1310-1311.
- Sparks, R.S.J., Cashman, K.V., 2017. Dynamic Magma Systems: Implications for Forecasting Volcanic Activity. *Elements* 13, 35-40.
- Spence, R., Kelman, I., Baxter, P., Zuccaro, G. and Petrazzuoli, S., 2005. Residential building and occupant vulnerability to tephra fall. *Natural Hazards and Earth System Science* 5, 477-494.
- Stevenson, J.A., Loughlin, S.C., Font, A., Fuller, G. W., MacLeod, A., Oliver, I.W., Jackson, B., Horwell, C.J., Thordarson, T., Dawson, I., 2013. UK monitoring and deposition of tephra from the May 2011 eruption of Grímsvötn, Iceland. *Journal of Applied Volcanology* 2, 3.
- Stone, J., Barclay, J., Simmons, P., Cole, P. D., Loughlin, S. C., Ramon, P., Mothes, P., 2014. Risk reduction through community-based monitoring: the vigías of Tungurahua, Ecuador. *Journal of Applied Volcanology* 3, 11.
- Surono, Jousset, P., Pallister, J., Boichu, M., Buongiorno, M. F., Budisantoso, A., Costa, F., Andreastuti, S., Prata, F., Schneider, D., Clar-

- isse, L., Humaida, H., Sumarti, S., Bignami, C., Griswold, J., Carn, S., Oppenheimer, C. & Lavigne, F., 2012. The 2010 explosive eruption of Java's Merapi volcano-A '100-year' event. *Journal of Volcanology and Geothermal Research* 241-242, 121-135.
- Suzuki, Y., Nagai, M., Maeno, F., Yasuda, A., Hokanishi, N., Shimano, T., Ichihara, M., Kaneko, T., Nakada, S., 2013. Precursory activity and evolution of the 2011 eruption of Shinmoe-dake in Kirishima volcano — insights from ash samples. *Earth Planets Space* 65, 591-607.
- Sword Daniels, V.L., Twigg, J., Loughlin, S.C., 2015. Time for change? Applying an inductive timeline tool for a retrospective study of disaster recovery in Montserrat, West Indies. *International Journal of Disaster Risk Reduction* 12, 125-133
- Sword-Daniels, V., 2011. Living with volcanic risk: The consequences of, and response to, ongoing volcanic ashfall from a social infrastructure systems perspective on Montserrat. *New Zealand Journal of Psychology* 40, 131-138.
- Sword-Daniels, V., Wilson, T. M., Sargeant, S., Rossetto, T., Twigg, J., Johnston, D. M., Loughlin, S. C., Cole, P. D., 2014. Consequences of long-term volcanic activity for essential services in Montserrat: challenges, adaptations and resilience. In: Wadge, G., Robertson, R.E.A., Voight, B., (eds.), *The Eruption of Soufrière Hills Volcano, Montserrat from 2000 to 2010*. Geological Society, London, UK, *Memoirs* 39, pp. 471-488.
- Thordarson, T. and Larsen, G. 2007. Volcanism in Iceland in historical time: Volcano types, eruption styles and eruptive history. *Journal of Geodynamics* 43, 118-152.
- Thordarson, T. and Self, S., 2003. Atmospheric and environmental effects of the 1783-1784 Laki eruption: a review and reassessment. *Journal of Geophysical Research: Atmospheres* (1984-2012) 108, AAC-7.
- Thorkelsson, B., 2012. The 2010 Eyjafjallajökull Eruption, Iceland: Report to ICAO. Icelandic Meteorological Office.
- Ulivieri G., Ripepe M., Marchetti, E., 2013. Infrasound reveals transition to oscillatory discharge regime during lava fountaining: implication for early warning. *Geophysical Research Letters* 40(12), 3008-3013.
- USGS, 2015. Volcano Hazards Program. Ash Cloud Simulations - What if Mount St. Helens Produced an Explosive Eruption Today?. https://volcanoes.usgs.gov/volcanoes/st_helens/monitoring_ash_cloud.html, [accessed 13 April, 2017].
- Venezky, D. Y. and Newhall, C.G., 2007, WOVODat design document; the schema, table descriptions, and create table statements for the database of worldwide volcanic unrest (WOVODat Version 1.0): U.S. Geological Survey Open File Report 2007-1117. <http://pubs.usgs.gov/of/2007/1117/>, [accessed 06 April, 2016].
- Villagran de Leon, J. C., 2012. Early warning principles and systems. In: Wisner, B., Gaillard, J. C. and Kelman, I. (Eds.), *The Routledge Handbook of Hazards and Disaster Risk Reduction*. Routledge, London, UK, 481-492.
- Voight, B., 1990. The 1985 Nevado del Ruiz volcano catastrophe: anatomy and retrospection. *Journal of Volcanology and Geothermal Research* 42, 151-188.
- Voight, B., Calvache, M. L., Minard V., Hall, L., Monsalve, M. L., 2013. The tragic 13 November 1985 eruption of Nevado del Ruiz Volcano, Colombia: the worst can happen. In: Bobrowsky, P. (ed.), *Encyclopedia of Natural Hazards*. Springer, Dordrecht, Netherlands.
- Voight, B., Sparks, R.S.J., Miller, A.D., Stewart, R.C., Hoblitt, R.P., Clarke, A., Ewart, J., Aspinall, W.P., Baptie, B., Calder, E.S., Cole, P., Druitt, T.H., Hartford, C., Herd, R.A., Jacksorn, P., Lejeune, A.M., Lockhart, A.B., Loughlin, S.C., Luckett, R., Lynch, L., Norton, G.E., Robertson, R., Watson, I.M., Watts, R. & Young, S.R., 1999. Magma Flow Instability and Cyclic Activity at Soufriere Hills Volcano, Montserrat, British West Indies. *Science* 283.5405, 1138-1142.
- Wadge, G. and Aspinall, W., 2014b. A review of volcanic hazard and risk assessments at the Soufrière Hills Volcano, Montserrat from 1997 to 2011. In: Wadge, G., Robertson, R., Voight, B. *The Eruption of Soufriere Hills Volcano, Montserrat, from 2000 to 2010*, Geological Society Memoirs, Vol. 39. Geological Society of London, London.
- Wadge, G., Voight, B., Sparks, R.S.J., Cole, P., Loughlin, S.C., 2014a. An overview of the eruption of Soufriere Hills Volcano from 2000-2010. In: Wadge, G., Robertson, R., Voight, B. *The Eruption of Soufriere Hills Volcano, Montserrat, from 2000 to 2010*. Geological Society Memoirs 39. Geological Society, London, 1-40.
- Wallace, K., Snedigar, S., Cameron, C. 2015. 'Is Ash Falling?', an online ashfall reporting tool in support of improved ashfall warnings and investigations of ashfall processes. *Journal of Applied Volcanology* 4, 8.
- Widiwijayanti, C., Costa, F., Nang, T. Z. W., Tan, K., Newhall, C., Ratdomopurbo, A., 2015. Recent development of WOVODat – The global volcano unrest database as a resource to improve eruption forecasts. In: G-EVER Promotion Team (Eds.). 2015 International Workshop on Earthquake and Volcanic Hazards and Risks in Asia-Pacific Region. G-EVER Consortium and Geological survey of Japan, National Institute of Advanced Industrial Science and Technology, Japan.
- Wilson, G., Wilson, T.M., Deligne, N.I., Cole, J.W., 2014. Volcanic hazard impacts to critical infrastructure: A review. *Journal of Volcanology and Geothermal Research* 286, 148-182.
- Wilson, T. M., Stewart, C., Bickerton, H., Baxter, P. J., Outes, V., Villarosa, G., Rovere, E., 2013. Impacts of the June 2011 Puyehue Cordón-Caulle volcanic complex eruption on urban infrastructure, agriculture and public health. *GNS Science Report* 2012/20.
- Wilson, T., Cole, J., Stewart, C., Cronin, S. and Johnston, D., 2011. Ash storms: impacts of wind-remobilised 889 volcanic ash on rural communities and agriculture following the 1991 Hudson eruption, southern 890 Patagonia, Chile. *Bulletin of Volcanology* 73(3), 223-239.
- Winson, A.E.G., Costa, F., Newhall, C.G., Woo, G., 2014. An analysis of the issuance of volcanic alert levels during volcanic crises. *Journal of Applied Volcanology* 3, 14.
- Wisner, B., Gaillard, J., Kelman, I. (eds.) 2012. *The Handbook of Hazards and Disaster Risk Reduction*, Routledge, Oxford.
- Witham, C., Oppenheimer, C., Horwell, C. J., 2005. Volcanic ash-leachates: a review and recommendations for sampling methods. *Journal of Volcanology and Geothermal Research* 41, 299-326.
- Woo, G. 2014. Cost-benefit analysis in volcanic risk, 289-300. In: Papale, P. (ed.) *Volcanic Hazards, Risks and Disasters*. Elsevier, 532pp.
- Zuccaro, G., Cacace, F., Spence, R. J. S. and Baxter, P. J., 2008. Impact of explosive eruption scenarios at Vesuvius. *Journal of Volcanology and Geothermal Research* 178, 416-453.
- Zuccaro, G., De Gregorio, D., Baxter, P., 2014. Human and Structural Vulnerability to Volcanic Processes, 261-188. In: Papale, P. *Volcanic Hazards, Risks and Disasters*. Elsevier, 532pp.

3.3 Geophysical risk: tsunamis

- Anunziato, A., 2015. The Inexpensive device for Sea level Measurements. *Science of Tsunami Hazards* 34(4), 199-211.
- ASTARTE, 2013. Assessment, Strategy And Risk Reduction for Tsunamis in Europe. <http://www.astarte-project.eu/index.php/astarte-home.html>, [accessed 15 April, 2017].
- Babeyko, A. Y., Hoehner, A., Sobolev, S. V., 2010. Source modeling and inversion with near real-time GPS: a GITWS perspective for Indonesia. *Natural Hazards and Earth Systems Sciences* 10(7), 1617-1627.
- Basili, R., Tiberti, M. M., Kastelic, V., Romano, F., Piatanesi, A., Selva, J., Lorito, S., 2013. Integrating geologic fault data into tsunami hazard studies. *Natural Hazards and Earth Systems Sciences* 13, 1025-1050.
- Blewitt, G., Kreemer, C., Hammond, W. C., Plag, H.-P., Stein, S., Okal, E., 2006. Rapid determination of earthquake magnitude using GPS for tsunami warning systems. *Geophysical Research Letters* 33(11), 11309.
- Burbidge D., Cummins P.R., Mleczo R., Thio, H.K., 2008. A Probabilistic Tsunami Hazard Assessment for Western Australia, *Pure and Applied Geophysics* 165, 2059.
- Davies G., Griffin J., Løvholt, F., Glymsdal, S., Harbitz, C., Thio, H.K., Lorito, S., Basili, R., Selva, J., Geist E., Baptista M.A. 2017. A global probabilistic tsunami hazard assessment from earthquake sources. Geological Society, London, Special Publications 456.
- EIDA, 2017. European Integrated Data Archive. ORFEUS EPOS. European Plate Observing System. <http://www.orfeus-eu.org/data/eida/>, [accessed 14 April 2017].
- EUREF, 2011. EUREF Permanent Network (EPN). http://www.euref.eu/euref_epn.html, [accessed 14 April 2017].
- Falck, C., Ramatschi, M., Subarya, C., Bartsch, M., Merx, A., Hoeberechts, J., Schmidt, G., 2010. Near real-time GPS applications for tsunami early warning systems. *Natural Hazards and Earth Systems Sciences* 10(2), 181-189.
- Geist, E. L., Parsons, T., 2006. Probabilistic Analysis of Tsunami Hazards. *Natural Hazards* 37, 277-314.
- Geller, R. J., 2011. Shake up time for Japanese seismology. *Nature* 472, 407-409.
- Grezio, A., Marzocchi, W., Sandri, L., Gasparini, P., 2010. A Bayesian procedure for Probabilistic Tsunami Hazard Assessment. *Natural Hazards* 53, 159-174.
- GTM, n.d. Global tsunami model. www.globaltsunamimodel.org, [accessed 16 April, 2017].
- Hoehner, A., Ge, M., Babeyko, A. Y., Sobolev, S.V., 2013. Instant tsunami early warning based on real-time GPS — Tohoku 2011 case study. *Natural Hazards and Earth Systems Sciences* 13(5), 1285-1292.
- Howe, B. M., Aucan, J., Tilmann, F., 2016. Submarine cable systems for future societal needs. *Eos, Transactions American Geophysical Union*, 97.
- IOC, 1998. Post-tsunami survey field guide, 1st edn, Intergovernmental Oceanographic Commission Manuals and Guides 37, Unesco.
- Kagan, Y. Y., Jackson, D. D., 2013. Tohoku Earthquake: A Surprise?. *Bulletin of the Seismological Society of America* 103, 1181-1194.
- Kanamori, H., 1977. The energy release in great earthquakes. *Journal of Geophysical Research*, 82, 2981-2987.
- Lomax, A., Michelini, A., 2009a. Mwpd: A Duration-Amplitude Procedure for Rapid Determination of Earthquake Magnitude and Tsunamigenic Potential from P Waveforms. *Geophysical Journal International* 176, 200-214.
- Lomax, A., Michelini, A., 2009b. Tsunami early warning using earthquake rupture duration. *Geophysical Research Letters* 36, L09306.
- Lorito S., Selva J., Basili, R., Romano F., Tiberti, M. M., Piatanesi, A., 2015. Probabilistic hazard for seismically induced tsunamis: accuracy and feasibility of inundation maps. *Geophysical Journal International* 200 (1), 574-588.
- Lorito, S., Romano, F., Lay, T., 2015. Tsunamigenic earthquakes (2004-2013): Source processes from data inversion. In: Meyers, R. (ed.), *Encyclopedia of Complexity and Systems Science*. Springer Science+Business Media, New York, USA.
- Lorito, S., Tiberti, M.M., Basili, R., Piatanesi, A., Valensise, G., 2008. Earthquake-generated tsunamis in the Mediterranean Sea: scenarios of potential threats to Southern Italy. *Journal of Geophysical Research* 113, B01301.
- Løvholt, F., Griffin, J., Salgado-Gálvez, M., 2015. Tsunami hazard and risk assessment at a global scale. In: Meyers, R. (ed.), *Encyclopedia of Complexity and Systems Science*. Springer Science+Business Media, New York, USA.
- Melgar, D., Allen, R. M., Riquelme, S., Geng, J., Bravo, F., Baez, J.C., Parra, H., Barrientos, S., Fang, P., Bock, Y., Bevis, M., Caccamise, D. J., Vigny, C., Moreno, M., Smalley, R., 2016. Local tsunami warnings: Perspectives from recent large events. *Geophysical Research Letters* 43 (3), 1109—1117.
- Michelini, A., Charalampakis, M., 2016. Working Group 2: Seismic and Geophysical measurements Report on intersessional activities. 13th session of the ICG for the Tsunami Early Warning and Mitigation System in the NEAM region (ICG/NEAMTWS-XIII), Bucharest, Romania, 26-28 September 2016. http://ioc-unesco.org/index.php?option=com_oe&task=viewDocumentRecord&docID=17804, [accessed 15 April, 2017].
- Miller, D., 1960. Giant waves in Lituya Bay Alaska. *USGS Professional Paper* 354-C, 51-83.
- Moustopolou, V., Nicol, A., Begg, J., Oncken, O., Moreno, M., 2015. Clusters of mega earthquakes on upper plate faults control the Eastern Mediterranean hazard. *Geophysical Research Letters* 42 (23), 10282-10289.
- Münch, U., Rudloff, A., Lauterjung, J., 2011. Postface 'The GITWS Project — results, summary and outlook'. *Natural Hazards and Earth System Sciences* 11, 765-769.
- NEAMWAVE12, 2012. NEAM Press Release 04 December 2012, Successful first test of Tsunami Warning System for the North Atlantic and Mediterranean. International Tsunami Information Center: http://itic.ioc-unesco.org/index.php?option=com_content&view=category&id=2105&Itemid=2421, [accessed 15 April, 2017].
- NEAMWAVE14, 2014. Exercise NEAMWave14: NEAMWave14 successfully undertaken. International Tsunami Information Center. http://itic.ioc-unesco.org/index.php?option=com_content&view=category&layout=blog&id=2161&Itemid=2609, [accessed 15 April, 2017].
- NGDC/WDS, n.d. National Geophysical Data Center / World Data Service: Global Historical Tsunami Database. National Geophysical Data Center, NOAA. https://www.ngdc.noaa.gov/hazard/tsu_db.shtml, [accessed 14 April, 2017].
- Ohta, Y., Kobayashi, T., Tushima, H., Miura, S., Hino, R., Takasu, T., Fujimoto, H., Iinuma, T., Tachibana, K., Demachi, T., Sato, T., Ohzo-

- no, M., Umino, N., 2012. Quasi real-time fault model estimation for near-field tsunamis forecasting based on RTK-GPS analysis: Application to the 2011 Tohoku-Oki earthquake (Mw 9.0). *Journal of Geophysical Research: Solid Earth*, 117, B02311.
- Omira, R., Matias, L., Baptista, M. A., 2016. Developing an Event-Tree Probabilistic Tsunami Inundation Model for NE Atlantic Coasts: Application to a Case Study. *Pure and Applied Geophysics* 173 (12), 3775-3794.
- Papadopoulos G. A., Gràcia, E., Urgeles, R., Sallares, V., De Martini, P. M., Pantosti, D., González, M., Yalciner, A. C., Mascle, J., Sakellariou, D., Salamon, A., Tinti, S., Karastathis, V., Fokaefs, A., Camerlenghi, A., Novikova, T., Papageorgiou, A., 2014. Historical and pre-historical tsunamis in the Mediterranean and its connected seas: Geological signatures, generation mechanisms and coastal impacts. *Marine Geology*, 354, 81-109.
- Papadopoulos, G. A. and Imamura, F., 2001. A proposal for a new tsunami intensity scale. In: *Proceedings of the International Tsunami Symposium 2001, Seattle, Session 5, paper 5-1, pp. 569-577.*
- Papadopoulos, G. A., 2015. *Tsunamis in the European-Mediterranean Region: From Historical Record to Risk Mitigation.* Elsevier, Amsterdam, Netherlands.
- Polet, J., Kanamori, H., 2009. Tsunami earthquakes. In: Meyers, A. (ed.), *Encyclopedia of Complexity and Systems Science.* Springer, New York, USA.
- Power, W., Wang, X., Lane, E. M., Gillibrand, P. A., 2013. A Probabilistic Tsunami Hazard Study of the Auckland Region, Part I: Propagation Modelling and Tsunami Hazard Assessment at the Shoreline. *Pure and Applied Geophysics* 170 (9-10), 1621-1634.
- Richter, C.F., 1935. An instrumental earthquake scale. *Bulletin of the Seismological Society of America* 25, 1-32.
- Rudloff, A., Lauterjung, J., Münch, U., Tinti, S., 2009. Preface 'The GITWS Project (German-Indonesian Tsunami Early Warning System). *Natural Hazards and Earth Systems Sciences* 9, 1381-1382.
- Schindelé, F., 1998. Tsunami warning in near field for the two large 1996 Peru earthquakes. In: *Proceedings of the International Conference on Tsunamis, Paris, France.*
- Schindelé, F., Gailler, A., Hébert, H., Loevenbruck, A., Gutierrez, E., Monnier, A., Roudil, P., Reymond, D., Rivera, L., 2015. Implementation and challenges of the tsunami warning system in the western Mediterranean. *Pure and Applied Geophysics*, 172 (3-4), 821 -833.
- Selva, J., Tonini, R., Molinari, I., Tiberti, M. M., Romano, F., Grezio, A., Melini, D., Piatanesi, A., Basili, R., Lorito, S., 2016. Quantification of source uncertainties in Seismic Probabilistic Tsunami Hazard Analysis (SPTHA). *Geophysics Journal International* 205(3), 1780-1803.
- Shearer, P., Bürgmann, R., 2010. Lessons learned from the 2004 Sumatra-Andaman megathrust rupture. *Annual Review of Earth and Planetary Sciences* 38, p. 103-131.
- Sieberg, A., (1927). *Geologische, physikalische und angewandte Erdbebenkunde.* Verlag von Gustav Fischer, Jena (in German).
- Sobolev, S.V. Babeyko, A.Y., Wang, R., Hoehner, A., Galas, R., Rothacher, M., Sein, D. V., Schröter, J., Lauterjung, J., Subarya, C., 2007. Tsunami early warning using GPS-Shield arrays. *Journal of Geophysical Research: Solid Earth* 112(B8), B08415.
- Song, Y.T., 2007. Detecting tsunami genesis and scales directly from coastal GPS stations. *Geophysics Research Letters* 34, L19602.
- Sørensen, M. B., Spada, M., Babeyko, A., Wiemer, S., Grünthal, G., 2012. Probabilistic tsunami hazard in the Mediterranean Sea. *Journal of Geophysics Research* 117, B01305.
- Synolakis, C. E., 2011. Tsunamis: When will we learn? *Newsweek Magazine.*
- Synolakis, C., Kánoğlu, U., 2015. The Fukushima accident was preventable. *Philosophical Transaction Royal Society AilTrans. R. Soc. A* 373, 20140379.
- Tinti, S., Armigliato, A., 2003. The use of scenarios to evaluate the tsunami impact in southern Italy. *Marine Geology* 199(3), 221-243.
- Tinti, S., Armigliato, A., Pagnoni, G., Zaniboni, F., 2005. Scenarios of giant tsunamis of tectonic origin in the Mediterranean. *ISET Journal of Earthquake Technology* 42(4), 171-188.
- Tonini, R., Armigliato, A., Pagnoni, G., Zaniboni, F., Tinti, S., 2011. Tsunami hazard for the city of Catania, eastern Sicily, Italy, assessed by means of Worst-case Credible Tsunami Scenario Analysis (WCTSA). *Natural Hazards and Earth Systems Sciences* 11, 1217-1232.
- TSUMAPS-NEAM, n.d. Probabilistic Tsunami Hazard maps for the NEAM region. <http://www.tsumaps-neam.eu>, [accessed 16 April, 2017].
- UNESCO/IOC, 2017. Sea level station monitoring facility. <http://ioc-sealevelmonitoring.org/>, [accessed 15 April 2017].
- UNISDR, 2013. *Global assessment report on disaster risk reduction — from shared risk to shared value: the business case for disaster risk reduction.* UNISDR, Geneva, Switzerland.
- UNISDR, 2015a. *Global assessment report on disaster risk reduction — making development sustainable: the future of disaster risk management.* UNISDR, Geneva, Switzerland.
- UNISDR, 2015b. *Sendai framework for disaster risk reduction 2015–2030.* United Nations International Strategy for Disaster Reduction. http://www.wcdrr.org/uploads/Sendai_Framework_for_Disaster_Risk_Reduction_2015-2030.pdf, [accessed 04 April 2016].
- UNISDR/CRED, 2016. *Tsunami Disaster Risk: Past impacts and projections.* United Nations Office for Disaster Risk Reduction (UNISDR), Centre for Research on the Epidemiology of Disasters (CRED). (http://www.preventionweb.net/files/50825_credtsunami08.pdf, [accessed 14 April 2017].
- Webcritech, n.d. <http://webcritech.jrc.ec.europa.eu>, [accessed 15 April 2017].